

Table 26.13
Cancer of the Thyroid (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2010-2012 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.01	0.03	0.59	0.06	Asian/ Pacific Islander	0	0.00	0.01	0.02	0.61	0.08
	10	0.01	0.03	0.09	0.60	0.06		10	0.01	0.02	0.08	0.62	0.09
	20	0.02	0.08	0.17	0.59	0.06		20	0.02	0.08	0.16	0.61	0.09
	30	0.06	0.15	0.27	0.58	0.06		30	0.06	0.14	0.26	0.60	0.09
	40	0.09	0.21	0.35	0.52	0.06		40	0.09	0.20	0.34	0.54	0.09
	50	0.12	0.27	0.39	0.44	0.06		50	0.11	0.25	0.38	0.47	0.09
	60	0.16	0.29	0.34	0.35	0.06		60	0.15	0.27	0.35	0.36	0.08
	70	0.15	0.21	-	0.22	0.05		70	0.14	0.22	-	0.24	0.07
80	0.09	-	-	0.10	0.04	80	0.10	-	-	0.13	0.05		
White	0	0.00	0.01	0.04	0.63	0.06	American Indian/ Alaska Native ^a	0	0.00	0.00	0.01	0.35	0.03
	10	0.01	0.03	0.10	0.64	0.06		10	0.00	0.01	0.05	0.35	0.03
	20	0.03	0.09	0.19	0.63	0.06		20	0.01	0.05	0.09	0.35	0.03
	30	0.07	0.17	0.29	0.61	0.06		30	0.04	0.08	0.16	0.35	0.03
	40	0.10	0.23	0.38	0.55	0.06		40	0.04	0.12	0.22	0.32	0.03
	50	0.13	0.28	0.41	0.47	0.06		50	0.08	0.19	0.29	0.29	0.03
	60	0.17	0.30	0.36	0.36	0.06		60	0.11	0.22	0.22	0.22	0.04
	70	0.16	0.22	-	0.23	0.05		70	0.13	0.13	-	0.13	0.04
80	0.09	-	-	0.10	0.04	80	0.01	-	-	0.01	0.02		
Black	0	0.00	0.00	0.01	0.27	0.03	Hispanic ^b	0	0.00	0.01	0.02	0.48	0.06
	10	0.00	0.01	0.03	0.28	0.04		10	0.01	0.02	0.05	0.48	0.06
	20	0.01	0.03	0.07	0.28	0.04		20	0.01	0.05	0.11	0.48	0.06
	30	0.02	0.07	0.12	0.27	0.04		30	0.04	0.10	0.19	0.47	0.06
	40	0.05	0.10	0.18	0.26	0.04		40	0.06	0.15	0.27	0.44	0.06
	50	0.06	0.14	0.21	0.22	0.04		50	0.09	0.21	0.32	0.38	0.06
	60	0.09	0.17	0.18	0.19	0.04		60	0.13	0.24	0.31	0.31	0.06
	70	0.09	0.11	-	0.12	0.03		70	0.13	0.21	-	0.21	0.05
80	0.03	-	-	0.04	0.03	80	0.10	-	-	0.11	0.04		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.