



Table 26.19
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

| State | Rate | SE | Rank(CI) ^d | PD |
|------------------|------|------|-----------------------|---------------------|
| TOTAL U.S. | 0.51 | 0.01 | | |
| High Five States | | | | |
| Rhode Island | 0.81 | 0.18 | 01(01,41) | 59.47 ^c |
| Utah | 0.66 | 0.12 | 02(01,41) | 30.58 ^c |
| Idaho | 0.65 | 0.14 | 03(01,42) | 27.47 ^c |
| Virginia | 0.64 | 0.06 | 04(01,27) | 25.67 ^c |
| Kansas | 0.64 | 0.10 | 05(01,40) | 25.41 ^c |
| Low Five States | | | | |
| Maryland | 0.42 | 0.06 | 38(08,42) | -17.61 ^c |
| Georgia | 0.40 | 0.05 | 39(15,42) | -21.83 ^c |
| Connecticut | 0.38 | 0.07 | 40(07,42) | -25.01 ^c |
| Louisiana | 0.37 | 0.06 | 41(11,42) | -26.73 ^c |
| Mississippi | 0.34 | 0.07 | 42(12,42) | -33.75 ^c |

| State | Rate | SE | Rank(CI) ^d | PD | State | Rate | SE | Rank(CI) ^d | PD |
|---------------|------|------|-----------------------|---------------------|----------------|------|------|-----------------------|---------------------|
| Alabama | 0.51 | 0.07 | 20(02,41) | 1.01 | Montana | - | - | -(-,-) | - |
| Alaska | - | - | -(-,-) | - | Nebraska | 0.63 | 0.12 | 06(01,41) | 23.18 ^c |
| Arizona | 0.56 | 0.06 | 10(01,37) | 10.69 | Nevada | 0.44 | 0.08 | 33(02,42) | -13.56 |
| Arkansas | 0.43 | 0.08 | 36(04,42) | -15.27 ^c | New Hampshire | 0.44 | 0.11 | 32(01,42) | -13.20 |
| California | 0.53 | 0.03 | 17(06,31) | 4.68 | New Jersey | 0.49 | 0.05 | 23(04,40) | -2.65 |
| Colorado | 0.59 | 0.08 | 07(01,39) | 16.27 ^c | New Mexico | 0.44 | 0.09 | 34(02,42) | -14.16 |
| Connecticut | 0.38 | 0.07 | 40(07,42) | -25.01 ^c | New York | 0.55 | 0.04 | 13(03,31) | 8.67 |
| Delaware | - | - | -(-,-) | - | North Carolina | 0.54 | 0.05 | 14(02,38) | 6.02 |
| D.C. | - | - | -(-,-) | - | North Dakota | - | - | -(-,-) | - |
| Florida | 0.45 | 0.03 | 31(16,40) | -11.83 | Ohio | 0.56 | 0.04 | 12(02,34) | 9.42 |
| Georgia | 0.40 | 0.05 | 39(15,42) | -21.83 ^c | Oklahoma | 0.56 | 0.08 | 11(01,41) | 9.61 |
| Hawaii | 0.51 | 0.12 | 19(01,42) | 1.17 | Oregon | 0.49 | 0.07 | 24(02,42) | -3.31 |
| Idaho | 0.65 | 0.14 | 03(01,42) | 27.47 ^c | Pennsylvania | 0.49 | 0.04 | 26(06,39) | -3.57 |
| Illinois | 0.43 | 0.04 | 35(13,42) | -14.87 | Rhode Island | 0.81 | 0.18 | 01(01,41) | 59.47 ^c |
| Indiana | 0.52 | 0.06 | 18(02,40) | 2.44 | South Carolina | 0.43 | 0.06 | 37(05,42) | -15.96 ^c |
| Iowa | 0.53 | 0.08 | 16(01,42) | 4.72 | South Dakota | - | - | -(-,-) | - |
| Kansas | 0.64 | 0.10 | 05(01,40) | 25.41 ^c | Tennessee | 0.48 | 0.06 | 29(03,42) | -5.13 |
| Kentucky | 0.48 | 0.07 | 28(02,42) | -5.00 | Texas | 0.58 | 0.04 | 08(02,28) | 14.06 |
| Louisiana | 0.37 | 0.06 | 41(11,42) | -26.73 ^c | Utah | 0.66 | 0.12 | 02(01,41) | 30.58 ^c |
| Maine | - | - | -(-,-) | - | Vermont | - | - | -(-,-) | - |
| Maryland | 0.42 | 0.06 | 38(08,42) | -17.61 ^c | Virginia | 0.64 | 0.06 | 04(01,27) | 25.67 ^c |
| Massachusetts | 0.57 | 0.06 | 09(01,37) | 12.20 | Washington | 0.51 | 0.06 | 21(03,40) | 0.03 |
| Michigan | 0.49 | 0.05 | 25(05,40) | -3.55 | West Virginia | 0.50 | 0.10 | 22(01,42) | -2.09 |
| Minnesota | 0.54 | 0.07 | 15(01,39) | 5.95 | Wisconsin | 0.47 | 0.06 | 30(04,42) | -7.23 |
| Mississippi | 0.34 | 0.07 | 42(12,42) | -33.75 ^c | Wyoming | - | - | -(-,-) | - |
| Missouri | 0.49 | 0.06 | 27(03,41) | -4.11 | | | | | |



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.