

Table 27.5
Cancer of the Urinary Bladder (Invasive and *In Situ*)
Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Year of Diagnosis:									
1975-2012	20.97	37.29	9.35	22.74	40.61	9.98	12.65	20.53	7.44
1975	19.32	34.37	8.88	20.23	36.39	9.17	10.56	16.34	6.26
1976	19.69	34.39	9.44	20.73	36.70	9.72	12.48	18.16	8.59
1977	18.99	33.40	8.88	20.01	35.61	9.18	13.45	22.67	7.06
1978	19.99	35.36	9.44	21.07	37.83	9.74	11.73	19.36	6.39
1979	19.98	35.35	9.39	21.30	38.19	9.84	10.35	15.61	6.54
1980	20.42	37.14	8.92	21.53	39.81	9.08	13.27	18.23	9.87
1981	20.69	36.38	9.73	21.94	38.77	10.29	11.60	20.72	5.48
1982	20.09	35.79	9.53	21.38	38.30	10.17	12.19	21.77	5.60
1983	20.04	35.99	9.01	21.31	38.69	9.41	12.17	18.70	7.57
1984	20.86	37.72	9.18	22.34	40.88	9.63	11.49	18.93	6.24
1985	20.67	36.37	9.87	22.11	39.30	10.36	13.06	20.33	8.14
1986	21.04	37.95	9.42	22.46	40.86	9.95	12.94	22.16	6.63
1987	21.66	39.52	9.24	23.17	42.82	9.58	13.22	21.78	7.41
1988	20.86	38.10	8.99	22.71	41.85	9.68	11.29	17.99	6.45
1989	21.02	37.41	9.63	22.72	40.91	10.17	11.37	17.61	7.16
1990	21.08	37.49	9.64	22.63	40.78	10.03	12.69	18.59	8.75
1991	20.92	37.63	9.33	22.71	41.26	9.89	12.81	19.44	8.50
1992	21.25	37.52	9.81	22.95	40.73	10.50	12.48	20.82	6.89
1993	21.25	37.65	9.72	22.85	40.52	10.41	14.32	24.76	7.85
1994	20.81	36.88	9.46	22.62	40.26	10.19	12.34	20.30	6.89
1995	20.64	36.23	9.56	22.45	39.51	10.29	11.85	18.23	7.39
1996	20.82	36.71	9.54	22.68	40.12	10.27	12.43	20.32	7.28
1997	21.09	37.15	9.59	22.98	40.68	10.23	12.76	20.15	8.21
1998	21.61	38.57	9.43	23.61	42.23	10.13	12.03	20.54	6.52
1999	21.78	38.25	9.96	23.66	41.71	10.57	14.59	22.48	9.55
2000	21.85	38.82	9.56	23.92	42.47	10.35	13.22	20.43	8.56
2001	21.78	38.58	9.66	24.09	42.56	10.63	11.72	18.72	7.23
2002	21.28	37.20	9.70	23.33	40.70	10.54	12.79	18.15	9.11
2003	21.59	37.81	9.77	23.78	41.56	10.62	12.97	21.26	7.48
2004	21.55	38.00	9.56	23.69	41.67	10.36	13.77	22.76	8.30
2005	21.98	38.83	9.63	24.08	42.61	10.29	14.02	23.07	8.11
2006	21.39	37.93	9.22	23.54	41.73	9.92	11.73	18.04	7.55
2007	21.93	39.24	9.02	24.20	43.06	9.84	13.07	21.48	7.70
2008	21.37	37.58	9.35	23.46	41.11	10.14	12.86	22.08	6.80
2009	20.88	37.00	8.80	22.81	40.25	9.44	12.95	22.26	6.93
2010	21.28	37.37	9.16	23.66	41.36	9.98	12.32	20.86	6.76
2011	20.30	35.55	8.75	22.62	39.34	9.64	12.49	21.17	6.98
2012	20.60	36.07	8.95	22.64	39.36	9.77	13.00	23.30	6.20

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.