

Table 27.10
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Survival Time																					
1-year	88.9	89.9	91.7	91.7	92.2	91.1	92.9	92.5	91.7	92.1	91.5	91.8	91.8	91.9	92.1	92.5	91.6	90.8	90.5	90.9	91.8
2-year	82.8	85.1	87.0	87.5	88.0	87.1	88.2	88.3	86.8	87.8	86.8	88.3	87.2	87.7	87.1	88.1	86.6	86.5	84.6	87.0	
3-year	79.5	81.9	84.2	85.2	85.5	84.9	85.2	84.7	84.2	84.6	85.3	86.1	84.9	85.1	84.3	85.4	83.2	83.8	82.1		
4-year	76.5	79.4	81.9	83.4	83.1	82.7	82.6	82.6	81.9	82.5	83.1	83.3	82.8	83.2	82.8	83.9	81.2	80.9			
5-year	74.3	77.4	80.1	81.6	82.9	81.0	79.9	80.2	79.2	81.0	82.3	82.0	81.0	82.0	81.9	82.4	79.7				
6-year	72.3	75.8	78.4	80.1	80.9	79.6	77.9	78.8	77.6	80.2	80.7	80.6	79.3	80.1	80.7	81.4					
7-year	70.0	74.4	76.6	78.2	79.2	78.4	76.7	77.8	76.5	78.3	78.6	79.7	77.8	77.5	79.8						
8-year	68.3	73.2	74.9	76.8	77.8	76.9	75.8	75.9	75.1	76.6	77.6	78.3	76.3	75.7							
9-year	66.4	71.4	73.7	75.3	75.6	75.5	74.2	74.8	74.4	75.3	76.8	76.6	75.1								
10-year	64.2	70.2	72.3	73.9	74.7	73.8	73.2	73.1	74.1	73.0	75.4	75.6									
11-year	62.2	68.9	71.3	72.7	74.0	72.5	72.2	72.0	73.2	70.8	74.6										
12-year	60.9	67.4	69.6	71.6	72.5	70.6	70.7	69.7	71.8	69.5											
13-year	59.5	66.4	68.3	70.8	70.6	69.6	69.9	68.8	70.5												
14-year	58.4	64.6	66.7	70.1	68.8	67.6	68.0	68.3													
15-year	57.1	63.1	65.3	68.4	68.0	67.3	66.3														
16-year	55.8	61.3	63.9	67.1	66.1	67.1															
17-year	54.6	60.3	63.1	65.8	64.9																
18-year	53.6	59.3	61.8	64.9																	
19-year	52.8	58.6	61.2																		
20-year	52.0	58.2	60.6																		

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).