

Table 27.15  
Cancer of the Urinary Bladder

SEER Incidence (Invasive and *In Situ*) and U.S. Mortality  
Age-Adjusted Rates and Trends<sup>a</sup>  
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas <sup>b</sup>			SEER 13 Areas <sup>b</sup>			SEER 9 Areas <sup>b</sup>		
	Rate 2008-2012			Trend 2003-2012 <sup>c</sup>			Trend 2003-2012 <sup>d</sup>		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	20.3	35.8	8.7	-0.8*	-0.9*	-1.1*	-0.6*	-0.7*	-1.0*
White	22.2	39.0	9.4	-0.6*	-0.8*	-1.0*	-0.5*	-0.7*	-1.0*
White Hispanic <sup>e</sup>	11.5	20.2	5.2	-0.2	-0.2	-0.4	-	-	-
White Non-Hispanic <sup>e</sup>	23.8	41.7	10.1	-0.5*	-0.6*	-0.9*	-	-	-
Black	12.6	21.4	6.9	0.2	0.5*	-2.7*	0.2*	0.4*	-2.7
Asian/Pacific Islander	9.0	15.8	4.0	-1.3*	-1.4	-0.1	-	-	-
Amer Ind/Alaska Nat <sup>f</sup>	8.5	15.6	3.4	1.5	1.7	4.2	-	-	-
Hispanic <sup>e</sup>	11.2	19.6	5.1	-0.2	-0.2	-0.4	-	-	-
U.S. Mortality <sup>g</sup>	Rate 2008-2012			Trend 2003-2012 <sup>h</sup>					
RACE/ETHNICITY	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	4.4	7.7	2.2	0.0	0.0	-0.5*			
White	4.6	8.1	2.2	0.3*	0.2*	-0.3*			
White Hispanic <sup>e</sup>	2.4	4.1	1.3	0.0	-0.3	0.1			
White Non-Hispanic <sup>e</sup>	4.8	8.4	2.3	0.3*	0.3*	-0.2*			
Black	3.5	5.3	2.5	-1.0*	-0.8*	-1.2*			
Asian/Pacific Islander	1.7	3.0	0.9	-0.7	0.1	-1.6*			
Amer Ind/Alaska Nat									
Total U.S.	2.0	3.3	1.1	1.4	2.0*	-0.4			
CHSDA Counties	2.3	4.2	1.1	2.5*	4.2*	-			
Non-CHSDA Counties	1.6	2.2	1.1	-0.4	-0.8	-			
Hispanic <sup>e</sup>	2.3	3.9	1.3	-0.2	-0.4	0.0			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.2.0, April 2015, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2003-2012 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 1992-2012.

d The 2003-2012 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2012.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2003-2012 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2003-2012 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2012.

\* The APC is significantly different from zero (p<.05).