



Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.40	0.02		
High Five States				
Maine	6.12 ^b	0.27	01(01,04)	38.87 ^c
Vermont	5.69	0.40	02(01,21)	29.25 ^c
Nevada	5.32 ^b	0.21	03(01,18)	20.80 ^c
Massachusetts	5.16 ^b	0.12	04(02,14)	17.15 ^c
Ohio	5.04 ^b	0.09	05(03,16)	14.36
Low Five States				
South Dakota	3.79	0.28	47(20,50)	-13.97
Alaska	3.76	0.44	48(05,51)	-14.61
Texas	3.70 ^b	0.06	49(43,50)	-16.01 ^c
Mississippi	3.59 ^b	0.15	50(38,50)	-18.41 ^c
Hawaii	2.56 ^b	0.18	51(50,51)	-41.79 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.93 ^b	0.12	43(29,50)	-10.87	Montana	4.79	0.28	17(02,42)	8.75
Alaska	3.76	0.44	48(05,51)	-14.61	Nebraska	4.05	0.20	39(18,50)	-8.04
Arizona	4.12	0.11	37(25,47)	-6.44	Nevada	5.32 ^b	0.21	03(01,18)	20.80 ^c
Arkansas	4.53	0.17	21(06,41)	2.85	New Hampshire	5.00	0.26	06(02,34)	13.47
California	3.94 ^b	0.05	42(37,48)	-10.60	New Jersey	4.90 ^b	0.10	08(04,21)	11.15
Colorado	3.85 ^b	0.13	45(32,50)	-12.61	New Mexico	3.81	0.19	46(28,50)	-13.39
Connecticut	4.83	0.15	13(03,28)	9.58	New York	4.43	0.06	27(18,34)	0.68
Delaware	4.69	0.30	18(02,46)	6.47	North Carolina	4.20	0.09	33(23,44)	-4.73
D.C.	4.00	0.38	41(04,50)	-9.06	North Dakota	4.45	0.33	26(02,50)	0.97
Florida	4.47	0.06	25(17,32)	1.45	Ohio	5.04 ^b	0.09	05(03,16)	14.36
Georgia	4.37	0.10	29(17,40)	-0.75	Oklahoma	4.48	0.15	24(09,41)	1.74
Hawaii	2.56 ^b	0.18	51(50,51)	-41.79 ^c	Oregon	4.85	0.15	10(03,27)	10.12
Idaho	4.18	0.23	36(10,50)	-5.14	Pennsylvania	4.81 ^b	0.08	15(05,22)	9.32
Illinois	4.40	0.08	28(18,37)	-0.07	Rhode Island	4.90	0.28	07(02,39)	11.23
Indiana	4.80	0.12	16(04,26)	9.00	South Carolina	4.29	0.13	30(17,44)	-2.60
Iowa	4.52	0.16	23(07,40)	2.56	South Dakota	3.79	0.28	47(20,50)	-13.97
Kansas	4.09	0.16	38(20,49)	-7.21	Tennessee	4.25	0.11	32(19,44)	-3.43
Kentucky	4.82	0.15	14(03,28)	9.42	Texas	3.70 ^b	0.06	49(43,50)	-16.01 ^c
Louisiana	4.18	0.14	34(20,47)	-5.10	Utah	3.91	0.20	44(22,50)	-11.13
Maine	6.12 ^b	0.27	01(01,04)	38.87 ^c	Vermont	5.69	0.40	02(01,21)	29.25 ^c
Maryland	4.89 ^b	0.13	09(03,24)	10.98	Virginia	4.18	0.10	35(22,45)	-5.12
Massachusetts	5.16 ^b	0.12	04(02,14)	17.15 ^c	Washington	4.84 ^b	0.12	11(04,25)	9.88
Michigan	4.84 ^b	0.09	12(04,22)	9.80	West Virginia	4.53	0.20	22(05,42)	2.83
Minnesota	4.04	0.12	40(27,48)	-8.30	Wisconsin	4.65	0.12	20(06,32)	5.63
Mississippi	3.59 ^b	0.15	50(38,50)	-18.41 ^c	Wyoming	4.68	0.41	19(02,50)	6.30
Missouri	4.27	0.11	31(19,43)	-3.08					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.