



Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.65	0.03		
High Five States				
Maine	10.88 ^b	0.56	01(01,03)	42.19 ^c
Vermont	9.61	0.80	02(01,33)	25.49 ^c
New Hampshire	8.91	0.55	03(01,32)	16.44 ^c
Massachusetts	8.88 ^b	0.24	04(02,19)	15.96 ^c
Ohio	8.80 ^b	0.18	05(02,18)	14.92
Low Five States				
Texas	6.40 ^b	0.13	47(41,50)	-16.41 ^c
Mississippi	6.33 ^b	0.34	48(30,50)	-17.27 ^c
D.C.	5.78	0.74	49(18,51)	-24.55 ^c
Alaska	5.74	0.82	50(11,51)	-25.03 ^c
Hawaii	4.73 ^b	0.37	51(49,51)	-38.15 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.10	0.27	36(20,48)	-7.29	Montana	8.66	0.59	07(01,41)	13.10
Alaska	5.74	0.82	50(11,51)	-25.03 ^c	Nebraska	7.03	0.41	40(16,49)	-8.15
Arizona	7.08	0.22	38(23,47)	-7.47	Nevada	8.67	0.42	06(02,29)	13.23
Arkansas	7.81	0.35	20(04,42)	2.09	New Hampshire	8.91	0.55	03(01,32)	16.44 ^c
California	6.76 ^b	0.10	42(35,47)	-11.69	New Jersey	8.59 ^b	0.21	09(02,22)	12.23
Colorado	6.60 ^b	0.27	44(29,50)	-13.71	New Mexico	6.60	0.38	45(22,50)	-13.75
Connecticut	8.46	0.32	16(02,29)	10.49	New York	7.81	0.14	21(16,31)	1.99
Delaware	7.49	0.61	30(02,50)	-2.18	North Carolina	7.32	0.20	31(21,44)	-4.41
D.C.	5.78	0.74	49(18,51)	-24.55 ^c	North Dakota	7.71	0.66	24(02,49)	0.67
Florida	7.66	0.12	26(19,33)	0.11	Ohio	8.80 ^b	0.18	05(02,18)	14.92
Georgia	7.66	0.23	27(14,39)	0.08	Oklahoma	7.62	0.31	28(08,43)	-0.51
Hawaii	4.73 ^b	0.37	51(49,51)	-38.15 ^c	Oregon	8.47	0.31	15(02,27)	10.69
Idaho	7.26	0.47	32(08,49)	-5.21	Pennsylvania	8.59 ^b	0.16	08(03,20)	12.28
Illinois	7.73	0.17	23(15,36)	1.06	Rhode Island	8.49	0.58	12(01,41)	10.94
Indiana	8.47	0.25	14(02,25)	10.72	South Carolina	7.07	0.28	39(20,48)	-7.58
Iowa	8.17	0.33	19(03,36)	6.72	South Dakota	6.50	0.56	46(15,51)	-15.08 ^c
Kansas	6.94	0.33	41(20,49)	-9.39	Tennessee	7.68	0.25	25(12,40)	0.29
Kentucky	8.52	0.31	11(02,27)	11.30	Texas	6.40 ^b	0.13	47(41,50)	-16.41 ^c
Louisiana	7.09	0.29	37(20,49)	-7.43	Utah	6.69	0.40	43(20,50)	-12.53
Maine	10.88 ^b	0.56	01(01,03)	42.19 ^c	Vermont	9.61	0.80	02(01,33)	25.49 ^c
Maryland	8.58	0.28	10(02,25)	12.16	Virginia	7.14	0.22	34(22,46)	-6.72
Massachusetts	8.88 ^b	0.24	04(02,19)	15.96 ^c	Washington	8.31	0.25	18(03,27)	8.57
Michigan	8.48 ^b	0.20	13(03,23)	10.81	West Virginia	7.49	0.40	29(07,47)	-2.17
Minnesota	7.13	0.25	35(22,47)	-6.87	Wisconsin	8.32	0.25	17(03,28)	8.69
Mississippi	6.33 ^b	0.34	48(30,50)	-17.27 ^c	Wyoming	7.78	0.80	22(02,50)	1.68
Missouri	7.22	0.23	33(21,46)	-5.72					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.