


Table 27.20
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2008-2012

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.19	0.02		
High Five States				
D.C.	3.02	0.43	01(01,46)	37.51 ^c
Delaware	2.84	0.31	02(01,39)	29.45 ^c
Vermont	2.81	0.37	03(01,45)	28.26 ^c
Massachusetts	2.76 ^b	0.11	04(01,15)	25.94 ^c
Maine	2.75	0.24	05(01,35)	25.57 ^c
Low Five States				
Colorado	1.84	0.12	47(28,50)	-16.24 ^c
Texas	1.84 ^b	0.06	48(36,50)	-16.30 ^c
Idaho	1.81	0.21	49(10,50)	-17.64 ^c
New Mexico	1.73	0.17	50(22,50)	-21.11 ^c
Hawaii	0.96 ^b	0.14	51(50,51)	-56.20 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	1.86	0.11	43(26,50)	-15.27 ^c	Montana	1.93	0.24	40(05,50)	-12.06
Alaska	2.06	0.46	35(01,51)	-6.23	Nebraska	2.03	0.19	37(07,50)	-7.59
Arizona	1.84 ^b	0.10	44(30,50)	-15.93 ^c	Nevada	2.70	0.20	06(01,32)	23.35 ^c
Arkansas	2.26	0.15	25(04,46)	3.19	New Hampshire	2.27	0.23	22(02,49)	3.66
California	1.96 ^b	0.04	39(32,46)	-10.50	New Jersey	2.54 ^b	0.09	08(02,27)	15.88 ^c
Colorado	1.84	0.12	47(28,50)	-16.24 ^c	New Mexico	1.73	0.17	50(22,50)	-21.11 ^c
Connecticut	2.44	0.14	13(02,37)	11.19	New York	2.26	0.06	24(12,35)	3.21
Delaware	2.84	0.31	02(01,39)	29.45 ^c	North Carolina	2.20	0.09	30(11,41)	0.46
D.C.	3.02	0.43	01(01,46)	37.51 ^c	North Dakota	2.09	0.30	33(01,50)	-4.64
Florida	2.08	0.05	34(23,42)	-5.06	Ohio	2.54 ^b	0.08	10(03,22)	15.65 ^c
Georgia	2.30	0.10	19(07,37)	4.70	Oklahoma	2.29	0.14	20(04,43)	4.53
Hawaii	0.96 ^b	0.14	51(50,51)	-56.20 ^c	Oregon	2.22	0.13	29(06,45)	1.04
Idaho	1.81	0.21	49(10,50)	-17.64 ^c	Pennsylvania	2.32	0.07	18(08,34)	5.95
Illinois	2.23	0.08	27(11,39)	1.51	Rhode Island	2.60	0.26	07(01,43)	18.62 ^c
Indiana	2.34	0.11	17(05,37)	6.88	South Carolina	2.45	0.13	11(02,35)	11.85
Iowa	2.04	0.14	36(13,50)	-7.17	South Dakota	1.90	0.26	41(04,50)	-13.13
Kansas	2.15	0.16	32(06,49)	-2.06	Tennessee	2.00	0.10	38(20,49)	-8.91
Kentucky	2.37	0.13	16(03,39)	7.95	Texas	1.84 ^b	0.06	48(36,50)	-16.30 ^c
Louisiana	2.29	0.13	21(05,43)	4.22	Utah	1.84	0.18	46(14,50)	-16.06 ^c
Maine	2.75	0.24	05(01,35)	25.57 ^c	Vermont	2.81	0.37	03(01,45)	28.26 ^c
Maryland	2.54	0.12	09(02,29)	15.76 ^c	Virginia	2.22	0.10	28(09,42)	1.11
Massachusetts	2.76 ^b	0.11	04(01,15)	25.94 ^c	Washington	2.37	0.11	15(04,36)	8.27
Michigan	2.40	0.09	14(05,32)	9.33	West Virginia	2.44	0.19	12(01,43)	11.48
Minnesota	1.90	0.11	42(27,50)	-13.28	Wisconsin	2.16	0.11	31(10,45)	-1.44
Mississippi	1.84	0.14	45(21,50)	-15.99 ^c	Wyoming	2.25	0.39	26(01,50)	2.81
Missouri	2.27	0.11	23(06,39)	3.61					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.