

Table 28.12  
Childhood Cancer (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2012  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <20 <sup>e</sup>	0 to <37 <sup>e</sup>	>=37 <sup>f</sup>	Complete <sup>g</sup>
<u>Race</u>	<u>Sex</u>										
All Races <sup>b</sup>	Both Sexes	65,893	56,052	50,529	45,384	38,463	33,067	234,055	326,769	+	+
	Males	34,508	29,895	26,046	23,607	19,837	16,567	122,721	167,928	+	+
	Females	31,385	26,157	24,483	21,777	18,626	16,500	111,334	158,841	+	+
White <sup>b</sup>	Both Sexes	53,100	46,757	42,637	37,801	32,859	29,117	194,105	275,292	+	+
	Males	27,958	25,069	21,916	19,706	17,019	14,676	102,124	141,990	+	+
	Females	25,142	21,688	20,721	18,095	15,840	14,441	91,981	133,302	+	+
Black <sup>b</sup>	Both Sexes	7,885	5,822	4,937	4,699	3,602	2,582	24,832	32,639	+	+
	Males	4,009	3,034	2,544	2,375	1,763	1,223	12,692	16,202	+	+
	Females	3,876	2,788	2,393	2,324	1,839	1,359	12,140	16,437	+	+
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	2,887	2,146	1,848	+	+	+	8,581	+	+	+
	Males	1,517	1,118	1,005	+	+	+	4,497	+	+	+
	Females	1,370	1,028	843	+	+	+	4,084	+	+	+
Hispanic <sup>d</sup>	Both Sexes	13,941	10,460	8,359	+	+	+	39,491	+	+	+
	Males	7,467	5,741	4,453	+	+	+	21,305	+	+	+
	Females	6,474	4,719	3,906	+	+	+	18,186	+	+	+

<sup>a</sup> US 2012 cancer prevalence counts are based on 2012 cancer prevalence proportions from the SEER registries and 1/1/2012 US population estimates based on the average of 2011 and 2012 population estimates from the US Bureau of the Census.

<sup>b c d</sup> Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 37 years for 1975-2012 SEER 9 data; 20 years for 1992-2012 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f g</sup> (f) Cases diagnosed more than 37 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <37 and >=37.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.