

Table 1.24
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity
Both Sexes

	All Races		White		Black	
	Rate ^b 2009-2013	APC ^c 2004-2013	Rate ^b 2009-2013	APC ^c 2004-2013	Rate ^b 2009-2013	APC ^c 2004-2013
All Sites	448.7	-0.9*	457.4	-0.9*	470.2	-1.3*
Breast	67.2	0.0	68.0	-0.1	86.7	-3.4*
Prostate ^f	59.0	-3.6*	58.8	-2.0*	71.2	0.5
Lung and Bronchus	57.3	-2.1*	56.3	-3.9*	65.0	-2.4*
Colon and Rectum	41.0	-2.9*	40.1	-3.0*	50.7	-2.9*
Melanoma of the Skin	21.8	0.9*	25.9	0.9*	18.1	1.6*
Urinary Bladder	20.1	-1.3*	22.0	-1.2*	15.5	-0.6
Non-Hodgkin Lymphoma	19.5	-0.5*	20.4	-0.6*	14.6	-0.7
Kidney and Renal Pelvis	15.6	1.1*	16.1	1.1*	14.0	2.7*
Thyroid	13.9	4.7*	14.7	4.7*	13.2	1.0*
Corpus and Uterus, NOS ^f	13.5	1.2*	14.3	0.4	12.6	-1.0*
Leukemia	13.5	0.4	13.7	0.9*	10.8	0.4
Pancreas	12.4	0.6*	12.4	0.7*	10.6	-2.5*
Oral Cavity and Pharynx	11.1	0.6*	11.6	0.9*	10.0	2.6*
Liver & IBD ^g	8.4	2.8*	7.5	3.6*	9.2	-2.0*
Stomach	7.4	-1.1*	7.1	-0.5*	8.3	4.5*
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2009-2013	APC ^c 2004-2013	Rate ^b 2009-2013	APC ^c 2004-2013	Rate ^b 2009-2013	APC ^c 2004-2013
All Sites	302.8	-1.1*	320.0	-1.3*	344.6	-1.3*
Breast	53.5	0.9*	43.7	-0.6	49.5	0.2
Lung and Bronchus	36.7	-1.5*	40.9	-1.3*	47.3	-4.7*
Colon and Rectum	34.8	-2.6*	38.9	-3.6*	34.7	-2.6*
Prostate ^f	30.1	-5.5*	28.4	-5.1*	29.7	-2.4*
Thyroid	13.6	4.6*	18.6	-0.7	17.7	-0.4
Liver & IBD ^g	13.5	-2.0*	12.9	0.0	15.4	0.8
Non-Hodgkin Lymphoma	13.5	-0.1	12.9	-0.2	12.9	2.0*
Corpus and Uterus, NOS ^f	11.1	2.3*	11.3	3.2	11.9	4.5*
Stomach	11.0	-3.6*	10.4	-1.0	11.4	2.3*
Pancreas	10.0	0.5*	9.7	-3.0	11.1	-0.3
Urinary Bladder	8.8	-1.8*	9.1	3.4*	11.1	-1.9*
Kidney and Renal Pelvis	8.1	1.4	8.8	5.0*	10.6	-2.2*
Leukemia	7.7	-0.1	8.2	0.2	10.6	0.3
Oral Cavity and Pharynx	7.7	-0.4	8.1	-1.8	6.7	0.3
Ovary ^{fh}	5.1	-0.8	5.6	-1.9	6.0	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

^d Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^e Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^h Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.25
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013
All Sites	504.5	-1.7*	All Sites	508.2	-1.7*	All Sites	571.8	-2.4*
Prostate	129.4	-4.0*	Prostate	121.9	-4.3*	Prostate	203.5	-3.8*
Lung and Bronchus	67.9	-2.7*	Lung and Bronchus	68.1	-2.7*	Lung and Bronchus	87.3	-3.1*
Colon and Rectum	47.1	-3.1*	Colon and Rectum	46.0	-3.3*	Colon and Rectum	59.2	-3.0*
Urinary Bladder	35.3	-1.4*	Urinary Bladder	38.5	-1.3*	Kidney and Renal Pelvis	25.0	1.1
Melanoma of the Skin	28.5	1.2*	Melanoma of the Skin	33.5	1.1*	Urinary Bladder	21.2	-0.8
Non-Hodgkin Lymphoma	23.7	-0.4*	Non-Hodgkin Lymphoma	24.8	-0.4*	Non-Hodgkin Lymphoma	17.6	-1.1
Kidney and Renal Pelvis	21.4	1.1*	Kidney and Renal Pelvis	21.9	1.1*	Pancreas	17.2	-0.3
Leukemia	17.3	0.3	Leukemia	18.2	0.2	Liver & IBD ^f	16.4	2.6*
Oral Cavity and Pharynx	16.7	0.7*	Oral Cavity and Pharynx	17.4	1.0*	Myeloma	15.7	0.7
Pancreas	14.1	0.6*	Pancreas	14.1	0.7*	Oral Cavity and Pharynx	14.4	-2.0*
Liver & IBD ^f	13.0	2.8*	Liver & IBD ^f	11.6	3.5*	Stomach	14.0	-3.4*
Stomach	10.0	-1.5*	Stomach	9.1	-1.1*	Leukemia	13.8	0.2
Myeloma	8.2	1.3*	Brain and ONS ^f	8.3	-0.6*	Larynx	8.7	-3.4*
Brain and ONS ^f	7.6	-0.5*	Esophagus	7.9	-0.8*	Esophagus	7.2	-4.5*
Esophagus	7.4	-1.4*	Myeloma	7.7	1.4*	Brain and ONS ^f	4.9	1.4*
Asian/Pacific Islander			American Indian/Alaska Native^d			Hispanic^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013
All Sites	317.3	-2.3*	All Sites	338.5	-1.6*	All Sites	382.9	-2.4*
Prostate	68.9	-5.7*	Prostate	63.9	-5.4*	Prostate	106.9	-5.0*
Lung and Bronchus	47.9	-2.1*	Lung and Bronchus	48.6	-2.1	Colon and Rectum	41.3	-3.0*
Colon and Rectum	41.2	-2.1*	Colon and Rectum	46.2	-0.1	Lung and Bronchus	36.6	-3.4*
Liver & IBD ^f	20.8	-1.7*	Kidney and Renal Pelvis	25.1	1.3	Non-Hodgkin Lymphoma	20.7	0.0
Non-Hodgkin Lymphoma	16.5	-0.1	Liver & IBD ^f	18.5	-0.4	Kidney and Renal Pelvis	20.1	0.1
Urinary Bladder	15.5	-1.8*	Non-Hodgkin Lymphoma	14.6	-0.6	Liver & IBD ^f	19.5	1.9*
Stomach	14.3	-4.0*	Urinary Bladder	14.4	-2.0	Urinary Bladder	19.4	-2.2*
Kidney and Renal Pelvis	11.7	2.0*	Stomach	13.5	-2.0	Stomach	13.5	-3.2*
Oral Cavity and Pharynx	11.1	0.5	Oral Cavity and Pharynx	13.2	4.6	Leukemia	12.8	0.2
Pancreas	11.0	0.3	Pancreas	10.9	-	Pancreas	11.8	-0.6
Leukemia	9.6	0.0	Leukemia	9.7	0.1	Oral Cavity and Pharynx	10.0	0.2
Thyroid	6.5	5.8*	Myeloma	5.2	-	Myeloma	7.5	0.9
Myeloma	4.7	0.7	Esophagus	5.0	-	Brain and ONS ^f	5.7	-1.7*
Brain and ONS ^f	4.3	0.8	Testis	4.9	-	Thyroid	5.0	3.9*
Esophagus	3.4	-4.8*	Melanoma of the Skin	4.2	-	Testis	5.0	2.3*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.26
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Females

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	409.9	-0.2		All Sites	422.5	-0.2		All Sites	400.5	-0.3	
Breast	125.0	0.2		Breast	128.0	0.1		Breast	125.2	0.6*	
Lung and Bronchus	49.4	-1.5*		Lung and Bronchus	51.8	-1.4*		Lung and Bronchus	50.0	-1.5*	
Colon and Rectum	36.0	-2.7*		Colon and Rectum	35.2	-2.7*		Colon and Rectum	44.8	-3.0*	
Corpus and Uterus, NOS	25.4	1.4*		Corpus and Uterus, NOS	26.0	1.1*		Corpus and Uterus, NOS	24.6	2.9*	
Thyroid	20.6	4.8*		Thyroid	21.9	4.8*		Pancreas	14.2	-0.8	
Melanoma of the Skin	16.9	0.4		Melanoma of the Skin	20.4	0.4		Kidney and Renal Pelvis	12.8	2.0*	
Non-Hodgkin Lymphoma	16.1	-0.7*		Non-Hodgkin Lymphoma	16.9	-0.8*		Thyroid	12.4	5.0*	
Ovary ^g	11.9	-1.7*		Ovary ^g	12.5	-1.8*		Non-Hodgkin Lymphoma	12.2	-0.3	
Pancreas	11.0	0.5*		Leukemia	11.1	0.4		Myeloma	11.5	1.2	
Kidney and Renal Pelvis	10.7	0.9		Kidney and Renal Pelvis	11.1	0.9		Ovary ^g	9.6	-1.1*	
Leukemia	10.5	0.4		Pancreas	10.9	0.7*		Cervix Uteri	8.9	-2.5*	
Urinary Bladder	8.6	-1.6*		Urinary Bladder	9.2	-1.5*		Leukemia	8.7	0.4	
Cervix Uteri	7.5	-1.7*		Cervix Uteri	7.5	-1.5*		Stomach	8.1	-1.7	
Oral Cavity and Pharynx	6.2	0.0		Oral Cavity and Pharynx	6.4	0.4		Urinary Bladder	6.9	-1.9*	
Brain and ONS ^f	5.4	-0.5		Brain and ONS ^f	6.0	-0.4		Oral Cavity and Pharynx	5.2	-2.1*	
Asian/Pacific Islander											
All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	296.7	-0.1		All Sites	310.7	-0.9*		All Sites	321.9	-0.2	
Breast	97.3	0.9*		Breast	81.2	-0.5		Breast	92.4	0.3	
Colon and Rectum	29.8	-3.1*		Colon and Rectum	36.8	-2.3*		Colon and Rectum	29.4	-2.2*	
Lung and Bronchus	28.5	-0.6		Lung and Bronchus	32.0	-5.1*		Lung and Bronchus	24.8	-1.3*	
Corpus and Uterus, NOS	20.3	2.3*		Corpus and Uterus, NOS	21.3	3.4		Corpus and Uterus, NOS	21.4	2.5*	
Thyroid	19.9	4.2*		Thyroid	13.6	7.0*		Thyroid	18.6	4.6*	
Non-Hodgkin Lymphoma	11.1	-0.2		Kidney and Renal Pelvis	13.3	-3.3		Non-Hodgkin Lymphoma	15.3	-0.7	
Ovary ^g	9.3	-0.7		Non-Hodgkin Lymphoma	11.6	-0.3		Kidney and Renal Pelvis	11.5	1.6*	
Pancreas	9.3	0.8		Ovary ^g	10.4	-1.9		Ovary ^g	10.6	-1.7*	
Stomach	8.5	-3.2*		Pancreas	8.6	-4.0		Pancreas	10.4	0.0	
Liver & IBD ^f	7.6	-2.5*		Liver & IBD ^f	8.1	0.1		Cervix Uteri	9.4	-4.5*	
Leukemia	6.2	-0.2		Stomach	8.0	0.5		Leukemia	8.8	0.2	
Cervix Uteri	6.2	-3.1*		Cervix Uteri	7.7	-0.8		Stomach	8.5	-1.0	
Kidney and Renal Pelvis	5.4	0.4		Leukemia	7.2	0.9		Liver & IBD ^f	7.4	2.3*	
Oral Cavity and Pharynx	4.9	-1.7		Oral Cavity and Pharynx	5.7	-		Urinary Bladder	5.0	-1.6*	
Urinary Bladder	3.9	-1.3		Myeloma	4.8	-		Myeloma	4.9	-0.5	
American Indian/Alaska Native ^d											
All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	296.7	-0.1		All Sites	310.7	-0.9*		All Sites	321.9	-0.2	
Breast	97.3	0.9*		Breast	81.2	-0.5		Breast	92.4	0.3	
Colon and Rectum	29.8	-3.1*		Colon and Rectum	36.8	-2.3*		Colon and Rectum	29.4	-2.2*	
Lung and Bronchus	28.5	-0.6		Lung and Bronchus	32.0	-5.1*		Lung and Bronchus	24.8	-1.3*	
Corpus and Uterus, NOS	20.3	2.3*		Corpus and Uterus, NOS	21.3	3.4		Corpus and Uterus, NOS	21.4	2.5*	
Thyroid	19.9	4.2*		Thyroid	13.6	7.0*		Thyroid	18.6	4.6*	
Non-Hodgkin Lymphoma	11.1	-0.2		Kidney and Renal Pelvis	13.3	-3.3		Non-Hodgkin Lymphoma	15.3	-0.7	
Ovary ^g	9.3	-0.7		Non-Hodgkin Lymphoma	11.6	-0.3		Kidney and Renal Pelvis	11.5	1.6*	
Pancreas	9.3	0.8		Ovary ^g	10.4	-1.9		Ovary ^g	10.6	-1.7*	
Stomach	8.5	-3.2*		Pancreas	8.6	-4.0		Pancreas	10.4	0.0	
Liver & IBD ^f	7.6	-2.5*		Liver & IBD ^f	8.1	0.1		Cervix Uteri	9.4	-4.5*	
Leukemia	6.2	-0.2		Stomach	8.0	0.5		Leukemia	8.8	0.2	
Cervix Uteri	6.2	-3.1*		Cervix Uteri	7.7	-0.8		Stomach	8.5	-1.0	
Kidney and Renal Pelvis	5.4	0.4		Leukemia	7.2	0.9		Liver & IBD ^f	7.4	2.3*	
Oral Cavity and Pharynx	4.9	-1.7		Oral Cavity and Pharynx	5.7	-		Urinary Bladder	5.0	-1.6*	
Urinary Bladder	3.9	-1.3		Myeloma	4.8	-		Myeloma	4.9	-0.5	
Hispanic ^e											
All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	296.7	-0.1		All Sites	310.7	-0.9*		All Sites	321.9	-0.2	
Breast	97.3	0.9*		Breast	81.2	-0.5		Breast	92.4	0.3	
Colon and Rectum	29.8	-3.1*		Colon and Rectum	36.8	-2.3*		Colon and Rectum	29.4	-2.2*	
Lung and Bronchus	28.5	-0.6		Lung and Bronchus	32.0	-5.1*		Lung and Bronchus	24.8	-1.3*	
Corpus and Uterus, NOS	20.3	2.3*		Corpus and Uterus, NOS	21.3	3.4		Corpus and Uterus, NOS	21.4	2.5*	
Thyroid	19.9	4.2*		Thyroid	13.6	7.0*		Thyroid	18.6	4.6*	
Non-Hodgkin Lymphoma	11.1	-0.2		Kidney and Renal Pelvis	13.3	-3.3		Non-Hodgkin Lymphoma	15.3	-0.7	
Ovary ^g	9.3	-0.7		Non-Hodgkin Lymphoma	11.6	-0.3		Kidney and Renal Pelvis	11.5	1.6*	
Pancreas	9.3	0.8		Ovary ^g	10.4	-1.9		Ovary ^g	10.6	-1.7*	
Stomach	8.5	-3.2*		Pancreas	8.6	-4.0		Pancreas	10.4	0.0	
Liver & IBD ^f	7.6	-2.5*		Liver & IBD ^f	8.1	0.1		Cervix Uteri	9.4	-4.5*	
Leukemia	6.2	-0.2		Stomach	8.0	0.5		Leukemia	8.8	0.2	
Cervix Uteri	6.2	-3.1*		Cervix Uteri	7.7	-0.8		Stomach	8.5	-1.0	
Kidney and Renal Pelvis	5.4	0.4		Leukemia	7.2	0.9		Liver & IBD ^f	7.4	2.3*	
Oral Cavity and Pharynx	4.9	-1.7		Oral Cavity and Pharynx	5.7	-		Urinary Bladder	5.0	-1.6*	
Urinary Bladder	3.9	-1.3		Myeloma	4.8	-		Myeloma	4.9	-0.5	

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

^g Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.27
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity
Both Sexes

All Races	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c	All Sites	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	168.5	-1.5*		All Sites	168.4	-1.4*		All Sites	197.9	-2.1*	
Lung and Bronchus	46.0	-2.3*		Lung and Bronchus	46.7	-2.2*		Lung and Bronchus	49.4	-2.7*	
Colon and Rectum	15.1	-2.5*		Colon and Rectum	14.7	-2.5*		Colon and Rectum	20.7	-3.0*	
Breast	12.0	-2.0*		Breast	11.6	-2.0*		Breast	17.4	-1.5*	
Pancreas	10.9	0.1		Pancreas	10.8	0.2*		Prostate ^f	15.9	-3.7*	
Prostate ^f	8.2	-2.7*		Prostate ^f	7.7	-2.5*		Pancreas	13.5	-0.4*	
Leukemia	6.9	-0.9*		Leukemia	7.1	-0.7*		Liver & IBD ^g	8.1	2.5*	
Liver & IBD ^g	6.1	2.8*		Non-Hodgkin Lymphoma	6.3	-2.3*		Myeloma	6.2	-0.8*	
Non-Hodgkin Lymphoma	6.0	-2.3*		Liver & IBD ^g	5.7	2.9*		Stomach	6.1	-3.1*	
Urinary Bladder	4.4	0.0		Brain and ONS ^g	4.7	0.3		Leukemia	5.8	-1.5*	
Brain and ONS ^g	4.3	0.2		Urinary Bladder	4.6	0.1*		Corpus and Uterus, NOS ^f	4.7	1.6*	
Ovary ^f	4.2	-2.5*		Ovary ^f	4.3	-2.5*		Non-Hodgkin Lymphoma	4.4	-1.9*	
Esophagus	4.1	-1.0*		Esophagus	4.3	-0.4*		Ovary ^f	3.8	-2.0*	
Kidney and Renal Pelvis	3.9	-0.8*		Kidney and Renal Pelvis	4.0	-0.7*		Esophagus	3.8	-4.4*	
Myeloma	3.3	-0.8*		Melanoma of the Skin	3.1	0.0		Kidney and Renal Pelvis	3.7	-1.4*	
Stomach	3.3	-2.5*		Myeloma	3.1	-0.8*		Urinary Bladder	3.6	-0.9*	
Asian/Pacific Islander				American Indian/Alaska Native ^d				Hispanic ^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013
All Sites	104.4	-1.3*		All Sites	153.6	-1.2*		All Sites	117.6	-1.2*	
Lung and Bronchus	24.2	-1.6*		Lung and Bronchus	37.8	-1.7*		Lung and Bronchus	19.7	-2.5*	
Colon and Rectum	10.5	-1.5*		Colon and Rectum	16.8	0.1		Colon and Rectum	12.0	-1.7*	
Liver & IBD ^g	9.7	-0.6		Liver & IBD ^g	10.2	1.7		Liver & IBD ^g	9.0	1.6*	
Pancreas	7.7	0.0		Pancreas	8.7	-0.9		Pancreas	8.6	-0.2	
Breast	6.3	-1.2*		Breast	8.2	-3.9*		Breast	8.0	-1.2*	
Stomach	5.7	-4.0*		Prostate ^f	7.7	-2.6*		Prostate ^f	6.8	-2.8*	
Non-Hodgkin Lymphoma	4.0	-1.4*		Kidney and Renal Pelvis	6.5	1.0		Stomach	5.4	-2.7*	
Leukemia	4.0	0.2		Stomach	5.3	-2.5		Non-Hodgkin Lymphoma	5.1	-1.2*	
Prostate ^f	3.6	-3.7*		Non-Hodgkin Lymphoma	4.5	-2.5*		Leukemia	4.9	0.0	
Ovary ^f	2.5	-1.4*		Leukemia	4.5	-2.1		Kidney and Renal Pelvis	3.5	-1.1*	
Brain and ONS ^g	2.0	0.8		Ovary ^f	3.7	-1.5		Ovary ^f	3.1	-1.8*	
Oral Cavity and Pharynx	1.9	-0.9		Esophagus	3.3	-2.8*		Brain and ONS ^g	2.9	0.3	
Kidney and Renal Pelvis	1.9	-0.3		Myeloma	2.9	-2.2		Myeloma	2.8	-0.7	
Urinary Bladder	1.7	0.1		Brain and ONS ^g	2.4	2.5		Urinary Bladder	2.3	-0.6	
Esophagus	1.7	-2.2*		Urinary Bladder	2.1	1.3		Esophagus	2.2	-1.5	

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.28
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

			Males					
All Races			White			Black		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013
All Sites	204.0	-1.8*	All Sites	202.9	-1.6*	All Sites	254.2	-2.7*
Lung and Bronchus	57.8	-2.9*	Lung and Bronchus	57.7	-2.9*	Lung and Bronchus	70.6	-3.4*
Prostate	20.7	-3.4*	Prostate	19.1	-3.3*	Prostate	44.2	-4.3*
Colon and Rectum	18.1	-2.6*	Colon and Rectum	17.6	-2.6*	Colon and Rectum	26.1	-2.7*
Pancreas	12.5	0.1	Pancreas	12.5	0.2*	Pancreas	15.0	-0.5
Leukemia	9.3	-0.9*	Leukemia	9.6	-0.8*	Liver & IBD ^f	12.8	2.7*
Liver & IBD ^f	9.1	2.9*	Liver & IBD ^f	8.3	3.1*	Stomach	8.8	-3.4*
Non-Hodgkin Lymphoma	7.7	-2.0*	Urinary Bladder	8.1	0.0	Leukemia	7.7	-1.7*
Urinary Bladder	7.7	-0.1*	Non-Hodgkin Lymphoma	8.1	-2.0*	Myeloma	7.5	-1.3*
Esophagus	7.4	-1.0*	Esophagus	7.7	-0.5*	Esophagus	6.6	-4.6*
Kidney and Renal Pelvis	5.7	-0.6*	Kidney and Renal Pelvis	5.8	-0.5*	Non-Hodgkin Lymphoma	5.7	-1.6*
Brain and ONS ^f	5.3	0.2	Brain and ONS ^f	5.7	0.3	Kidney and Renal Pelvis	5.5	-1.4*
Stomach	4.5	-2.8*	Melanoma of the Skin	4.6	0.2	Urinary Bladder	5.4	-0.6
Myeloma	4.2	-0.8*	Myeloma	4.0	-0.7*	Oral Cavity and Pharynx	5.0	-3.4*
Melanoma of the Skin	4.1	0.0	Stomach	3.9	-2.8*	Larynx	3.5	-4.0*
Oral Cavity and Pharynx	3.8	-0.4	Oral Cavity and Pharynx	3.7	0.2	Brain and ONS ^f	3.1	0.7
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013
All Sites	125.6	-1.6*	All Sites	183.0	-1.3*	All Sites	145.0	-1.6*
Lung and Bronchus	32.7	-2.4*	Lung and Bronchus	46.6	-1.7	Lung and Bronchus	28.3	-3.3*
Liver & IBD ^f	14.4	-0.4	Prostate	19.1	-3.3*	Prostate	17.1	-3.3*
Colon and Rectum	12.6	-1.4*	Colon and Rectum	18.9	-1.1	Colon and Rectum	15.4	-1.3*
Prostate	9.1	-3.6*	Liver & IBD ^f	14.3	3.7*	Liver & IBD ^f	13.0	1.5*
Pancreas	8.3	-0.4	Pancreas	9.6	-1.9	Pancreas	9.6	-0.4
Stomach	7.5	-4.3*	Kidney and Renal Pelvis	9.0	-0.5	Stomach	7.1	-3.0*
Leukemia	5.1	-0.4	Stomach	7.4	-4.2	Non-Hodgkin Lymphoma	6.3	-1.0*
Non-Hodgkin Lymphoma	4.9	-1.5*	Leukemia	6.2	-1.7	Leukemia	6.2	-0.2
Urinary Bladder	2.9	0.7	Non-Hodgkin Lymphoma	5.7	0.1	Kidney and Renal Pelvis	4.9	-1.9*
Esophagus	2.9	-2.0	Esophagus	5.2	-4.2	Esophagus	4.1	-1.4
Oral Cavity and Pharynx	2.9	-0.9	Urinary Bladder	3.8	1.2	Urinary Bladder	4.0	-0.6
Kidney and Renal Pelvis	2.8	0.5	Myeloma	3.4	-3.0	Myeloma	3.5	-0.4
Brain and ONS ^f	2.4	0.5	Oral Cavity and Pharynx	3.3	-2.2	Brain and ONS ^f	3.4	0.1
Myeloma	2.2	-0.1	Brain and ONS ^f	3.0	1.9	Oral Cavity and Pharynx	2.4	-1.3*
Soft Tissue including Heart	1.1	0.6	Soft Tissue including Heart	1.7	-	Larynx	1.6	-3.0*

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.29
Age-Adjusted U.S. Death Rates and Trends for the Top 15 Cancer Sites^a by Race/Ethnicity

Females

All Races	Rate ^b		APC ^c	All Races	Rate ^b		APC ^c	All Races	Rate ^b		APC ^c
	2009-2013	2004-2013			2009-2013	2004-2013			2009-2013	2004-2013	
All Sites	143.4	-1.4*		All Sites	143.6	-1.3*		All Sites	163.8	-1.6*	
Lung and Bronchus	37.0	-1.6*		Lung and Bronchus	38.3	-1.5*		Lung and Bronchus	35.3	-1.9*	
Breast	21.5	-1.8*		Breast	21.0	-1.8*		Breast	29.6	-1.4*	
Colon and Rectum	12.7	-2.7*		Colon and Rectum	12.3	-2.6*		Colon and Rectum	17.1	-3.5*	
Pancreas	9.5	0.1		Pancreas	9.4	0.2		Pancreas	12.2	-0.3	
Ovary	7.5	-2.2*		Ovary	7.8	-2.2*		Corpus and Uterus, NOS	7.9	1.9*	
Leukemia	5.2	-1.0*		Leukemia	5.3	-0.9*		Ovary	6.5	-1.8*	
Non-Hodgkin Lymphoma	4.7	-2.8*		Non-Hodgkin Lymphoma	4.9	-2.8*		Myeloma	5.4	-0.5	
Corpus and Uterus, NOS	4.5	1.3*		Corpus and Uterus, NOS	4.1	1.1*		Leukemia	4.6	-1.4*	
Liver & IBD ^f	3.6	2.2*		Brain and ONS ^f	3.8	0.3		Liver & IBD ^f	4.4	2.0*	
Brain and ONS ^f	3.5	0.1		Liver & IBD ^f	3.4	2.3*		Stomach	4.2	-3.0*	
Myeloma	2.7	-1.1*		Kidney and Renal Pelvis	2.5	-1.3*		Cervix Uteri	3.9	-1.9*	
Kidney and Renal Pelvis	2.5	-1.4*		Myeloma	2.4	-1.2*		Non-Hodgkin Lymphoma	3.5	-2.0*	
Stomach	2.4	-2.4*		Urinary Bladder	2.2	-0.4*		Urinary Bladder	2.5	-1.7*	
Cervix Uteri	2.3	-0.7*		Cervix Uteri	2.1	-0.4		Kidney and Renal Pelvis	2.5	-1.5*	
Urinary Bladder	2.2	-0.6*		Stomach	2.1	-2.4*		Brain and ONS ^f	2.1	0.1	
Asian/Pacific Islander				American Indian/Alaska Native ^d				Hispanic ^e			
	Rate ^b	APC ^c		Rate ^b	APC ^c		Rate ^b	APC ^c			
	2009-2013	2004-2013		2009-2013	2004-2013		2009-2013	2004-2013			
All Sites	89.6	-1.0*		All Sites	132.3	-1.4*		All Sites	98.5	-0.9*	
Lung and Bronchus	18.1	-0.4		Lung and Bronchus	31.3	-1.6*		Breast	14.5	-1.0*	
Breast	11.2	-1.3*		Colon and Rectum	15.1	1.0		Lung and Bronchus	13.5	-1.4*	
Colon and Rectum	9.0	-1.7*		Breast	14.7	-3.7*		Colon and Rectum	9.4	-2.4*	
Pancreas	7.3	0.4		Pancreas	8.0	-0.9		Pancreas	7.8	-0.1	
Liver & IBD ^f	6.0	-0.9		Liver & IBD ^f	6.8	-1.6		Liver & IBD ^f	5.7	1.4*	
Ovary	4.5	-1.3*		Ovary	6.7	-1.3		Ovary	5.5	-1.7*	
Stomach	4.4	-3.7*		Kidney and Renal Pelvis	4.5	2.6		Non-Hodgkin Lymphoma	4.2	-1.4*	
Non-Hodgkin Lymphoma	3.3	-1.2*		Stomach	3.8	-0.8		Stomach	4.1	-2.5*	
Leukemia	3.1	0.9		Corpus and Uterus, NOS	3.6	3.7		Leukemia	4.0	0.4	
Corpus and Uterus, NOS	2.9	2.8*		Non-Hodgkin Lymphoma	3.5	-5.4*		Corpus and Uterus, NOS	3.6	2.6*	
Cervix Uteri	1.8	-1.7		Leukemia	3.3	-2.0		Cervix Uteri	2.6	-2.8*	
Brain and ONS ^f	1.7	1.1		Cervix Uteri	3.2	-1.3		Brain and ONS ^f	2.4	0.6	
Myeloma	1.3	-2.0		Myeloma	2.6	-1.1		Kidney and Renal Pelvis	2.3	-0.4	
Oral Cavity and Pharynx	1.2	-1.2		Esophagus	1.8	-		Myeloma	2.3	-1.2	
Kidney and Renal Pelvis	1.2	-1.5		Brain and ONS ^f	1.8	-		Urinary Bladder	1.2	-1.4	

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^a Top 15 cancer sites selected based on 2009-2013 age-adjusted rates for the race/ethnic group.

^b Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.


^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

Figure 1.12

SEER Cancer Incidence and US Death Rates, 2009-2013 By Cancer Site and Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG) and US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.


^a Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.13

SEER Incidence 2004-2013 Males by Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.


^a Incidence rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.14

SEER Incidence 2004-2013 Females by Race/Ethnicity



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.


^a Incidence rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 1.15

US Mortality 2004-2013 Males by Race/Ethnicity



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

^a Mortality rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Figure 1.16

US Mortality 2004-2013 Females by Race/Ethnicity



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

^a Mortality rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Table 2.15
All Cancer Sites (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	448.7	504.5	409.9	-1.1*	-2.0*	-0.3	-1.0*	-1.9*	-0.2*
White	457.4	508.2	422.5	-1.1*	-2.0*	-0.3	-1.1*	-1.9*	-0.2*
White Hispanic ^e	354.8	391.3	333.8	-1.2*	-2.7*	0.1*	-	-	-
White Non-Hispanic ^e	477.2	529.2	440.8	-1.0*	-1.9*	-0.2	-	-	-
Black	470.2	571.8	400.5	-1.5*	-2.6*	-0.6	-1.4*	-2.3*	0.0
Asian/Pacific Islander	302.8	317.3	296.7	-1.1*	-2.4*	0.1	-	-	-
Amer Ind/Alaska Nat ^f	320.0	338.5	310.7	0.0	-0.5	0.5*	-	-	-
Hispanic ^e	344.6	382.9	321.9	-1.1*	-2.3*	0.1	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	168.5	204.0	143.4	-1.5*	-1.8*	-1.4*			
White	168.4	202.9	143.6	-1.4*	-1.7*	-1.4*			
White Hispanic ^e	124.2	152.8	104.1	-1.1*	-1.3*	-0.9*			
White Non-Hispanic ^e	172.4	207.2	147.3	-1.4*	-1.6*	-1.3*			
Black	197.9	254.2	163.8	-2.1*	-2.7*	-1.6*			
Asian/Pacific Islander	104.4	125.6	89.6	-1.4*	-1.6*	-1.0*			
Amer Ind/Alaska Nat									
Total U.S.	114.4	136.6	98.5	-2.0*	-1.4*	-2.1*			
CHSDA Counties	153.6	183.0	132.3	-1.1*	-0.3*	-1.3*			
Non-CHSDA Counties	70.4	85.1	60.1	-2.8*	-3.3*	-2.5*			
Hispanic ^e	117.6	145.0	98.5	-1.2*	-1.6*	-1.0*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 2.4

SEER Incidence and US Death Rates^a All Cancer Sites, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 3.15
Cancer of the Brain And Other Nervous System (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	6.4	7.6	5.4	-0.3*	-0.4*	-0.2	-0.2*	-0.4*	-0.1
White	7.1	8.3	6.0	-0.1	-0.3*	-0.1	-0.1	-0.3*	0.0
White Hispanic ^e	5.2	6.0	4.5	-1.6*	-0.2	-0.9*	-	-	-
White Non-Hispanic ^e	7.6	9.0	6.4	0.1	-0.1	0.2*	-	-	-
Black	4.1	4.9	3.6	0.0	0.2	-0.1	0.2	0.0	0.5*
Asian/Pacific Islander	3.7	4.3	3.1	-0.2	-0.2	-0.3	-	-	-
Amer Ind/Alaska Nat ^f	3.2	3.9	2.5	-5.5	0.1	-	-	-	-
Hispanic ^e	4.9	5.7	4.3	-1.8*	-2.4*	-0.9*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	4.3	5.3	3.5	0.0	0.0	0.0			
White	4.7	5.7	3.8	0.1	0.1	0.1			
White Hispanic ^e	3.1	3.6	2.6	0.2	0.1	0.2			
White Non-Hispanic ^e	4.9	6.0	4.0	0.3	0.2	0.2			
Black	2.5	3.1	2.1	-0.7*	-0.5*	-0.8*			
Asian/Pacific Islander	2.0	2.4	1.7	0.2	0.5	-0.1			
Amer Ind/Alaska Nat									
Total U.S.	2.0	2.3	1.7	0.2	-0.1	0.6			
CHSDA Counties	2.4	3.0	1.8	1.2	1.6	1.0			
Non-CHSDA Counties	1.5	1.5	1.5	-1.4	-2.8	-0.1			
Hispanic ^e	2.9	3.4	2.4	0.0	-0.1	0.1			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 3.2

SEER Incidence and US Death Rates^a Cancer of the Brain and Other Nervous System, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 4.18
Cancer of the Female Breast (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	125.0	0.2	0.3
White	128.0	0.0	0.1
White Hispanic ^e	96.1	0.2	-
White Non-Hispanic ^e	134.3	0.1	-
Black	125.2	0.2*	0.3*
Asian/Pacific Islander	97.3	1.1*	-
Amer Ind/Alaska Nat ^f	81.2	-0.1	-
Hispanic ^e	92.4	0.1	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Females	Females	
All Races	21.5	-1.9*	
White	21.0	-1.9*	
White Hispanic ^e	15.3	-1.0*	
White Non-Hispanic ^e	21.5	-1.8*	
Black	29.6	-1.5*	
Asian/Pacific Islander	11.2	-1.0*	
Amer Ind/Alaska Nat			
Total U.S.	11.1	-3.9*	
CHSDA Counties	14.7	-0.7	
Non-CHSDA Counties	7.1	-3.4*	
Hispanic ^e	14.5	-0.8	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.

^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Table 4.19
Cancer of the Female Breast (*In Situ*)

SEER Incidence
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013 Rate per 100,000 persons	Trend 2004-2013 ^c AAPC (%)	Trend 2004-2013 ^d AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	31.5	0.4	0.6*
White	31.5	0.2	0.4
White Hispanic ^e	21.9	1.8*	-
White Non-Hispanic ^e	33.8	0.0	-
Black	31.0	1.6*	1.5*
Asian/Pacific Islander	30.5	1.3*	-
Amer Ind/Alaska Nat ^f	16.5	1.6	-
Hispanic ^e	21.3	1.7*	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.


^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC is significantly different from zero ($p < .05$).

Figure 4.4

SEER Incidence and US Death Rates^a Cancer of the Female Breast

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 5.11
Cancer of the Cervix Uteri (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	7.5	-2.4*	-0.9*
White	7.5	-2.1*	-0.7
White Hispanic ^e	9.9	-4.0*	-
White Non-Hispanic ^e	7.0	-1.8*	-
Black	8.9	-3.5*	-3.6*
Asian/Pacific Islander	6.2	-4.1*	-
Amer Ind/Alaska Nat ^f	7.7	-1.2	-
Hispanic ^e	9.4	-4.1*	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Females	Females	
All Races	2.3	-0.8*	
White	2.1	-0.5*	
White Hispanic ^e	2.8	-2.7*	
White Non-Hispanic ^e	2.0	-0.1	
Black	3.9	-2.0*	
Asian/Pacific Islander	1.8	-3.5*	
Amer Ind/Alaska Nat			
Total U.S.	2.1	-3.2*	
CHSDA Counties	3.2	-2.3*	
Non-CHSDA Counties	0.9	-4.3*	
Hispanic ^e	2.6	-2.7*	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.


h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 5.3

SEER Incidence and US Death Rates^a Cancer of the Cervix Uteri

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 6.21
Cancer of the Colon and Rectum (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	41.0	47.1	36.0	-3.2*	-3.3*	-3.1*	-3.2*	-3.4*	-3.2*
White	40.1	46.0	35.2	-3.3*	-3.6*	-3.1*	-3.4*	-3.7*	-3.2*
White Hispanic ^e	35.8	42.9	30.2	-2.3*	-2.3*	-2.4*	-	-	-
White Non-Hispanic ^e	41.0	46.7	36.2	-3.4*	-3.7*	-3.1*	-	-	-
Black	50.7	59.2	44.8	-3.4*	-3.0*	-3.7*	-3.5*	-2.8*	-4.0*
Asian/Pacific Islander	34.8	41.2	29.8	-2.6*	-2.5*	-3.4*	-	-	-
Amer Ind/Alaska Nat ^f	40.9	46.2	36.8	-0.4	-0.3	-0.4	-	-	-
Hispanic ^e	34.7	41.3	29.4	-2.3*	-2.4*	-2.4*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	15.1	18.1	12.7	-2.7*	-2.7*	-2.8*			
White	14.7	17.6	12.3	-2.6*	-2.7*	-2.8*			
White Hispanic ^e	12.7	16.3	10.0	-1.5*	-1.2*	-2.0*			
White Non-Hispanic ^e	14.9	17.7	12.5	-2.6*	-2.8*	-2.8*			
Black	20.7	26.1	17.1	-3.0*	-2.7*	-3.4*			
Asian/Pacific Islander	10.5	12.6	9.0	-1.8*	-1.9*	-1.7*			
Amer Ind/Alaska Nat									
Total U.S.	12.1	13.9	10.8	-1.4*	-2.0*	-0.8			
CHSDA Counties	16.8	18.9	15.1	0.1	-1.0	0.0			
Non-CHSDA Counties	6.9	8.5	5.9	-1.9*	-2.0*	-1.5*			
Hispanic ^e	12.0	15.4	9.4	-1.7*	-1.4*	-2.2*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Table 6.22
Cancer of the Colon (Invasive) and Cancer of the Rectum (Invasive)

SEER Incidence
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

Cancer of the Colon <u>SEER Incidence</u>	SEER 18 Areas ^b Rate 2009-2013 Rate per 100,000 persons			SEER 13 Areas ^b Trend 2004-2013 ^c AAPC (%)			SEER 9 Areas ^b Trend 2004-2013 ^d AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
	RACE/ETHNICITY								
All Races	29.0	32.1	26.5	-3.5*	-3.6*	-3.4*	-3.3*	-3.7*	-3.2*
White	28.4	31.4	26.0	-3.6*	-3.9*	-3.4*	-3.6*	-4.0*	-3.1*
White Hispanic ^e	24.7	28.4	21.8	-2.5*	-2.5*	-2.5*	-	-	-
White Non-Hispanic ^e	29.1	32.0	26.6	-3.6*	-3.9*	-3.1*	-	-	-
Black	37.9	43.1	34.3	-3.5*	-3.4*	-4.0*	-3.6*	-3.4*	-4.3*
Asian/Pacific Islander	22.9	25.8	20.6	-3.0*	-3.0*	-3.5*	-	-	-
Amer Ind/Alaska Nat ^f	28.8	31.1	27.3	-0.7	-0.9	-0.5	-	-	-
Hispanic ^e	23.9	27.3	21.3	-2.4*	-2.5*	-2.5*	-	-	-
Cancer of the Rectum <u>SEER Incidence</u>	SEER 18 Areas ^b Rate 2009-2013 Rate per 100,000 persons			SEER 13 Areas ^b Trend 2004-2013 ^c AAPC (%)			SEER 9 Areas ^b Trend 2004-2013 ^d AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
	RACE/ETHNICITY								
All Races	12.0	15.0	9.5	-2.2*	-2.4*	-2.5*	-2.3*	-2.6*	-2.2*
White	11.7	14.6	9.3	-2.5*	-2.8*	-2.7*	-2.7*	-3.1*	-2.5*
White Hispanic ^e	11.1	14.5	8.3	-2.6*	-1.1*	-3.1	-	-	-
White Non-Hispanic ^e	11.9	14.8	9.5	-2.6*	-3.0*	-2.4*	-	-	-
Black	12.9	16.1	10.5	-2.8*	-1.0*	-3.4*	-2.5*	-2.1	-3.4*
Asian/Pacific Islander	11.9	15.4	9.2	-1.8*	-1.3*	-2.8*	-	-	-
Amer Ind/Alaska Nat ^f	12.1	15.1	9.5	0.3	0.9	-0.4	-	-	-
Hispanic ^e	10.8	14.0	8.2	-2.6*	-1.1*	-1.0*	-	-	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.


^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC is significantly different from zero (p<.05).

Figure 6.2

SEER Incidence and US Death Rates^a Cancer of the Colon and Rectum, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 7.11
Cancer of the Corpus And Uterus, NOS (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	25.4	1.1	1.0
White	26.0	1.1*	0.7
White Hispanic ^e	22.2	3.1*	-
White Non-Hispanic ^e	26.6	0.7	-
Black	24.6	2.3*	2.8*
Asian/Pacific Islander	20.3	1.8*	-
Amer Ind/Alaska Nat ^f	21.3	2.7*	-
Hispanic ^e	21.4	3.0*	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Females	Females	
All Races	4.5	1.2*	
White	4.1	1.1*	
White Hispanic ^e	3.8	2.3*	
White Non-Hispanic ^e	4.1	1.0*	
Black	7.9	1.7*	
Asian/Pacific Islander	2.9	1.9*	
Amer Ind/Alaska Nat			
Total U.S.	2.5	0.0	
CHSDA Counties	3.6	0.9	
Non-CHSDA Counties	1.2	-1.3	
Hispanic ^e	3.6	2.1*	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.

^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 7.3

SEER Incidence and US Death Rates^a Cancer of the Corpus and Uterus, NOS

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 8.15
Cancer of the Esophagus (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	4.3	7.4	1.7	-0.5*	-1.4*	-1.3*	-1.4*	-1.4*	-2.2*
White	4.5	7.9	1.7	-0.9	-0.9	-0.6*	-0.6	-0.7	-0.1
White Hispanic ^e	2.9	5.2	1.1	-1.1*	-1.3*	-1.2	-	-	-
White Non-Hispanic ^e	4.8	8.3	1.8	-0.8	-0.7	-0.4	-	-	-
Black	4.4	7.2	2.5	-4.6*	-4.7*	-4.3*	-4.5*	-4.8*	-3.3*
Asian/Pacific Islander	2.0	3.4	0.9	-2.0*	-2.3*	-0.7	-	-	-
Amer Ind/Alaska Nat ^f	3.0	5.0	1.4	-1.4	-2.5	-	-	-	-
Hispanic ^e	2.8	4.9	1.0	-1.1*	-1.3*	-1.3	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	4.1	7.4	1.5	-1.0*	-1.0*	-1.7*			
White	4.3	7.7	1.5	-0.4*	-0.5*	-1.1*			
White Hispanic ^e	2.3	4.3	0.8	-2.3*	-2.4*	-2.4*			
White Non-Hispanic ^e	4.5	8.0	1.6	-0.2	-0.4*	-1.0*			
Black	3.8	6.6	1.9	-4.5*	-4.6*	-4.1*			
Asian/Pacific Islander	1.7	2.9	0.8	-1.9*	-1.9*	-1.5*			
Amer Ind/Alaska Nat									
Total U.S.	2.4	4.0	1.2	-3.5*	-3.6*	0.2			
CHSDA Counties	3.3	5.2	1.8	-2.9	0.0	2.6			
Non-CHSDA Counties	1.4	2.6	-	-2.2*	-1.6	-4.5			
Hispanic ^e	2.2	4.1	0.8	-2.4*	-1.0*	-2.5*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 8.2

SEER Incidence and US Death Rates^a Cancer of the Esophagus, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 9.15
Hodgkin Lymphoma

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	2.6	3.0	2.3	-1.3*	-1.0*	-0.2	-1.2*	-0.8	-1.5
White	2.8	3.1	2.5	-1.5*	-0.1	-1.9	-1.4*	-0.7	-1.8
White Hispanic ^e	2.3	2.6	2.0	0.0	-0.5	0.5	-	-	-
White Non-Hispanic ^e	3.0	3.4	2.7	-1.7*	0.1	-2.1*	-	-	-
Black	2.6	3.0	2.2	0.8*	1.0*	0.4	1.1*	0.5	1.8*
Asian/Pacific Islander	1.3	1.5	1.2	-0.2	1.3	2.3*	-	-	-
Amer Ind/Alaska Nat ^f	1.2	1.5	1.0	-	-	-	-	-	-
Hispanic ^e	2.2	2.5	1.9	0.0	-0.5	0.5	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	0.4	0.5	0.3	-2.8*	-2.4*	-2.9*			
White	0.4	0.5	0.3	-2.7*	-2.3*	-2.8*			
White Hispanic ^e	0.4	0.6	0.3	-2.2*	-2.3*	-2.2*			
White Non-Hispanic ^e	0.4	0.5	0.3	-2.7*	-2.6*	-2.8*			
Black	0.3	0.4	0.3	-2.9*	-3.3*	-2.5*			
Asian/Pacific Islander	0.1	0.2	0.1	-2.1*	-2.7*	-			
Amer Ind/Alaska Nat									
Total U.S.	0.1	-	-	-3.0	-	-			
CHSDA Counties	-	-	-	-1.0	-	-			
Non-CHSDA Counties	-	-	-	-	-	-			
Hispanic ^e	0.4	0.6	0.3	-2.4*	-2.4*	-2.4*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.


- Statistic not shown. Rate based on less than 16 cases for the time interval.
- Trend based on less than 10 cases for at least one year within the time interval.
- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.
- The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.
- The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.
- ^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.
- ^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.
- ^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.
- ^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.
- ^g US Mortality Files, National Center for Health Statistics, CDC.
- ^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.
- * The APC is significantly different from zero (p<.05).

Figure 9.2

SEER Incidence and US Death Rates^a Hodgkin Lymphoma, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Joinpoint could not fit a regression line for American Indian/Alaska Native Incidence due to insufficient data.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 10.13
Kaposi Sarcoma

SEER Incidence
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	0.5	1.0	0.1	-2.7*	-2.7*	-4.3*	-1.8*	-1.8*	-3.9*
White	0.4	0.8	0.1	-3.3*	-3.4*	-4.4*	-2.5*	-2.6*	-4.8*
White Hispanic ^e	0.7	1.2	0.2	-4.6*	-4.2*	-3.7*	-	-	-
White Non-Hispanic ^e	0.4	0.7	0.1	-3.3*	-3.4*	-5.3*	-	-	-
Black	1.1	2.2	0.1	-1.6	-1.6	1.2	-0.1	-0.4	-
Asian/Pacific Islander	0.2	0.4	-	-1.2	-0.4	-	-	-	-
Amer Ind/Alaska Nat ^f	-	-	-	-	-	-	-	-	-
Hispanic ^e	0.7	1.2	0.2	-4.7*	-4.4*	-3.1	-	-	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC is significantly different from zero (p<.05).

Table 11.15
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	15.6	21.4	10.7	1.1*	1.2*	0.6	1.1*	1.7	0.5
White	16.1	21.9	11.1	1.0*	1.0*	0.7	1.0*	1.1*	0.5
White Hispanic ^e	16.0	21.0	12.0	0.8	0.3	2.6*	-	-	-
White Non-Hispanic ^e	16.2	22.2	10.9	1.1*	1.2*	0.4	-	-	-
Black	18.1	25.0	12.8	1.1	2.4*	2.2*	2.7*	2.8*	2.6*
Asian/Pacific Islander	8.1	11.7	5.4	2.5*	2.5*	0.7	-	-	-
Amer Ind/Alaska Nat ^f	18.6	25.1	13.3	1.5*	1.6*	1.2	-	-	-
Hispanic ^e	15.4	20.1	11.5	0.9	0.3	2.5*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	3.9	5.7	2.5	-0.9*	-0.8*	-0.8*			
White	4.0	5.8	2.5	-0.5*	-0.7*	-0.8*			
White Hispanic ^e	3.7	5.2	2.5	0.0	-0.2	0.1			
White Non-Hispanic ^e	4.0	5.9	2.5	-0.5*	-0.4*	-0.8*			
Black	3.7	5.5	2.5	-0.9*	-1.3*	-0.8*			
Asian/Pacific Islander	1.9	2.8	1.2	0.8	0.2	-0.4			
Amer Ind/Alaska Nat									
Total U.S.	4.3	5.7	3.1	-0.8*	-1.0	-0.9			
CHSDA Counties	6.5	9.0	4.5	0.2	-0.1	0.0			
Non-CHSDA Counties	1.7	2.1	1.4	-2.2*	-1.8	-2.2			
Hispanic ^e	3.5	4.9	2.3	-1.1*	-1.7*	0.0			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 11.2

SEER Incidence and US Death Rates^a Cancer of the Kidney and Renal Pelvis, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 12.15
Cancer of the Larynx (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	3.2	5.6	1.1	-2.6*	-2.6*	-3.1*	-2.4*	-2.4*	-2.7*
White	3.2	5.6	1.2	-2.4*	-1.8*	-2.9*	-2.3*	-2.4*	-2.4*
White Hispanic ^e	2.3	4.6	0.5	-1.7*	-1.6*	-3.2*	-	-	-
White Non-Hispanic ^e	3.4	5.8	1.3	-2.4*	-2.5*	-2.7*	-	-	-
Black	4.6	8.7	1.6	-3.4*	-3.4*	-3.4*	-3.0*	-2.8*	-3.1*
Asian/Pacific Islander	1.2	2.3	0.3	-3.1*	-3.0*	-2.4	-	-	-
Amer Ind/Alaska Nat ^f	2.1	3.9	-	-0.1	0.5	-	-	-	-
Hispanic ^e	2.2	4.4	0.5	-1.7*	-1.6*	-3.0*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	1.1	1.9	0.4	-2.2*	-2.5*	-2.1*			
White	1.0	1.8	0.4	-1.9*	-2.2*	-1.8*			
White Hispanic ^e	0.8	1.7	0.2	-2.5*	-2.7*	-2.1*			
White Non-Hispanic ^e	1.0	1.8	0.4	-1.8*	-2.2*	-1.6*			
Black	1.8	3.5	0.6	-3.4*	-3.4*	-3.1*			
Asian/Pacific Islander	0.4	0.8	0.1	-2.0*	-1.3*	-4.6*			
Amer Ind/Alaska Nat									
Total U.S.	0.7	1.2	0.3	-2.8*	-2.5	-			
CHSDA Counties	0.8	1.5	-	-2.8*	-2.2	-			
Non-CHSDA Counties	0.5	0.9	-	-	-	-			
Hispanic ^e	0.8	1.6	0.2	-2.5*	-2.7*	-2.2*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 12.2

SEER Incidence and US Death Rates^a Cancer of the Larynx, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 13.23
Leukemia

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	13.5	17.3	10.5	0.2*	0.1	0.3*	0.3*	0.1*	0.3*
White	14.3	18.2	11.1	0.4*	0.1	0.5*	0.4*	0.2*	0.4*
White Hispanic ^e	11.0	13.3	9.2	0.5*	0.3	0.7*	-	-	-
White Non-Hispanic ^e	14.6	18.7	11.2	0.4*	0.2*	0.5*	-	-	-
Black	10.8	13.8	8.7	0.2	0.2	0.2	0.0	0.0	0.0
Asian/Pacific Islander	7.7	9.6	6.2	-0.2	-0.2	-0.1	-	-	-
Amer Ind/Alaska Nat ^f	8.2	9.7	7.2	1.7	2.2	1.7	-	-	-
Hispanic ^e	10.6	12.8	8.8	0.5*	0.3	0.7*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	6.9	9.3	5.2	-1.0*	-1.0*	-1.2*			
White	7.1	9.6	5.3	-0.9*	-0.7*	-1.1*			
White Hispanic ^e	5.2	6.5	4.2	0.2	-0.3	-0.4*			
White Non-Hispanic ^e	7.2	9.8	5.3	-0.8*	-0.7*	-1.1*			
Black	5.8	7.7	4.6	-1.2*	-1.2*	-1.2*			
Asian/Pacific Islander	4.0	5.1	3.1	-0.7*	-0.6*	-0.7			
Amer Ind/Alaska Nat									
Total U.S.	3.6	4.8	2.7	-1.2*	-1.0	-1.5*			
CHSDA Counties	4.5	6.2	3.3	-0.3	0.1	-0.9			
Non-CHSDA Counties	2.5	3.3	2.0	-2.1*	-2.0	-2.1			
Hispanic ^e	4.9	6.2	4.0	0.1	-0.4*	-0.5*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.


h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 13.3

SEER Incidence and US Death Rates^a Leukemia, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 14.15
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	8.4	13.0	4.4	2.7*	2.3*	2.4*	3.0*	2.8*	2.7*
White	7.5	11.6	3.9	3.8*	3.8*	4.0*	4.3*	4.0*	4.0*
White Hispanic ^e	13.5	20.4	7.7	3.0*	2.9*	2.7*	-	-	-
White Non-Hispanic ^e	6.5	10.1	3.3	3.9*	3.9*	3.8*	-	-	-
Black	10.0	16.4	4.9	3.7*	2.5*	2.7*	3.1*	3.8*	2.8*
Asian/Pacific Islander	13.5	20.8	7.6	-2.1	-1.8	-2.4	-	-	-
Amer Ind/Alaska Nat ^f	12.9	18.5	8.1	2.5*	3.1*	1.2	-	-	-
Hispanic ^e	12.9	19.5	7.4	3.0*	2.9*	2.7*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	6.1	9.1	3.6	2.6*	2.8*	2.1*			
White	5.7	8.3	3.4	2.9*	3.0*	2.3*			
White Hispanic ^e	9.6	13.8	6.1	2.0*	2.0*	1.7*			
White Non-Hispanic ^e	5.3	7.8	3.2	2.9*	2.9*	2.4*			
Black	8.1	12.8	4.4	2.5*	2.4*	1.1*			
Asian/Pacific Islander	9.7	14.4	6.0	-0.8*	-0.8*	-0.7*			
Amer Ind/Alaska Nat									
Total U.S.	7.1	9.8	4.8	1.8*	1.8*	1.3			
CHSDA Counties	10.2	14.3	6.8	2.3*	2.6*	1.5			
Non-CHSDA Counties	3.6	4.8	2.5	2.0*	1.8	2.8*			
Hispanic ^e	9.0	13.0	5.7	1.9*	1.8*	1.6*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 14.2

SEER Incidence and US Death Rates^a Cancer of the Liver and Intrahepatic Bile Duct, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 15.21
Lung and Bronchus (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
RACE/ETHNICITY									
All Races	57.3	67.9	49.4	-2.1*	-2.9*	-1.5*	-1.8*	-2.6*	-1.2*
White	58.8	68.1	51.8	-2.1*	-2.8*	-1.5*	-1.8*	-2.6*	-1.2*
White Hispanic ^e	30.7	37.9	25.6	-1.3*	-2.1*	-0.4*	-	-	-
White Non-Hispanic ^e	63.2	72.7	56.2	-2.0*	-2.7*	-1.4*	-	-	-
Black	65.0	87.3	50.0	-2.7*	-2.7*	-1.7*	-1.9*	-2.8*	-1.3*
Asian/Pacific Islander	36.7	47.9	28.5	-1.4*	-2.2*	0.2	-	-	-
Amer Ind/Alaska Nat ^f	38.9	48.6	32.0	-0.9*	-1.6*	-3.5	-	-	-
Hispanic ^e	29.7	36.6	24.8	-1.3*	-2.1*	-0.4*	-	-	-
U.S. Mortality ^g									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
RACE/ETHNICITY									
All Races	46.0	57.8	37.0	-2.3*	-2.9*	-1.6*			
White	46.7	57.7	38.3	-2.1*	-2.8*	-1.4*			
White Hispanic ^e	20.7	29.7	14.2	-2.3*	-3.0*	-1.2*			
White Non-Hispanic ^e	49.1	60.2	40.6	-2.0*	-2.7*	-1.4*			
Black	49.4	70.6	35.3	-2.7*	-3.4*	-1.9*			
Asian/Pacific Islander	24.2	32.7	18.1	-1.2*	-2.2*	-0.4*			
Amer Ind/Alaska Nat									
Total U.S.	30.3	37.5	24.9	-1.2*	-2.1*	-2.3*			
CHSDA Counties	37.8	46.6	31.3	-0.3	-1.1*	-1.2			
Non-CHSDA Counties	21.7	27.1	17.5	-2.3*	-3.3*	-2.2*			
Hispanic ^e	19.7	28.3	13.5	-2.5*	-3.3*	-1.4*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.


h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 15.4

SEER Incidence and US Death Rates^a Cancer of the Lung and Bronchus, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 16.15
Melanoma of the Skin (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	21.8	28.5	16.9	1.0*	1.2*	0.6	1.4*	1.6*	1.0*
White	25.9	33.5	20.4	1.1*	1.2*	0.7	1.4*	1.6*	1.0*
White Hispanic ^e	4.6	4.9	4.5	0.9*	1.5*	0.4	-	-	-
White Non-Hispanic ^e	30.7	38.9	24.6	1.4*	1.6*	1.2*	-	-	-
Black	1.1	1.1	1.0	0.9	-2.9*	0.1	-0.1	-0.5	0.0
Asian/Pacific Islander	1.3	1.6	1.1	0.4	1.1	-0.4	-	-	-
Amer Ind/Alaska Nat ^f	4.4	4.2	4.6	2.1*	1.6	2.2	-	-	-
Hispanic ^e	4.5	4.8	4.4	0.7*	1.3*	0.3	-	-	-
<u>U.S. Mortality^g</u>									
RACE/ETHNICITY	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	2.7	4.1	1.7	0.0	0.3*	-0.5*			
White	3.1	4.6	2.0	0.2*	0.4*	-0.3*			
White Hispanic ^e	0.8	1.1	0.6	-0.2	-0.1	-0.4			
White Non-Hispanic ^e	3.4	5.0	2.1	0.4*	0.6*	-0.1			
Black	0.4	0.5	0.4	-1.1*	-0.1	-1.8*			
Asian/Pacific Islander	0.3	0.4	0.3	-1.1	-1.5	-0.8			
Amer Ind/Alaska Nat									
Total U.S.	0.7	1.0	0.5	0.6	1.5	-0.2			
CHSDA Counties	1.0	1.5	0.6	1.1	2.6	-			
Non-CHSDA Counties	0.5	-	-	1.2	-	-			
Hispanic ^e	0.8	1.0	0.6	-0.3	-0.2	-0.4			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.

^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 16.2

SEER Incidence and US Death Rates^a Melanoma of the Skin, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 17.11
Mesothelioma

SEER Incidence
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	1.0	1.8	0.4	-1.2*	-1.6*	-0.2	-1.1*	-1.6*	-0.2
White	1.1	2.0	0.5	-0.8*	-1.3*	0.3	-0.7*	-1.3*	0.7*
White Hispanic ^e	0.9	1.5	0.4	-5.3*	-2.3*	-1.4	-	-	-
White Non-Hispanic ^e	1.1	2.0	0.5	-0.7*	-1.2*	0.4	-	-	-
Black	0.5	1.0	0.3	-3.5*	-3.7*	-2.4	-4.7*	-4.5*	-
Asian/Pacific Islander	0.3	0.6	0.2	-1.3	-1.1	-0.9	-	-	-
Amer Ind/Alaska Nat ^f	-	-	-	-	-	-	-	-	-
Hispanic ^e	0.8	1.5	0.4	-5.4*	-2.2*	-1.5	-	-	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC is significantly different from zero (p<.05).

Table 18.15
Myeloma

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	6.5	8.2	5.2	0.8	0.8*	0.4*	0.8*	0.9*	0.7*
White	5.9	7.7	4.5	0.7*	0.8*	0.3	0.8*	0.9*	0.5*
White Hispanic ^e	6.1	7.7	4.9	0.2	0.4	-0.1	-	-	-
White Non-Hispanic ^e	5.9	7.7	4.4	0.7*	0.8*	0.3	-	-	-
Black	13.2	15.7	11.5	0.8*	1.1*	0.4	0.9*	0.9*	0.9*
Asian/Pacific Islander	3.7	4.7	3.0	0.6	0.7	0.6	-	-	-
Amer Ind/Alaska Nat ^f	5.0	5.2	4.8	0.2	-0.2	0.2	-	-	-
Hispanic ^e	6.0	7.5	4.9	0.2	0.4	-0.1	-	-	-
<u>U.S. Mortality^g</u>									
RACE/ETHNICITY	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	3.3	4.2	2.7	-0.9*	-1.0*	-1.0*			
White	3.1	4.0	2.4	-0.9*	-0.8*	-1.2*			
White Hispanic ^e	2.9	3.6	2.4	-0.6*	-0.6	-0.7*			
White Non-Hispanic ^e	3.1	4.0	2.4	-0.8*	-0.9*	-1.2*			
Black	6.2	7.5	5.4	-0.8	-1.4*	-0.5			
Asian/Pacific Islander	1.6	2.2	1.3	-0.8*	-0.3	-1.2*			
Amer Ind/Alaska Nat									
Total U.S.	2.2	2.5	2.0	-2.1*	-2.0*	-2.5*			
CHSDA Counties	2.9	3.4	2.6	-1.8*	-1.3	-2.3			
Non-CHSDA Counties	1.4	1.6	1.2	-1.7	-2.7	-			
Hispanic ^e	2.8	3.5	2.3	-0.7*	-0.7*	-0.8*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.


^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 18.2

SEER Incidence and US Death Rates^a Myeloma, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 19.19
Non-Hodgkin Lymphoma

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	19.5	23.7	16.1	-0.6	-0.6	-0.9*	-0.6	-0.6	-0.9*
White	20.4	24.8	16.9	-0.7	-0.5	-0.9*	-0.6*	-0.5	-1.0*
White Hispanic ^e	18.3	21.5	15.8	-0.6	0.2	0.7*	-	-	-
White Non-Hispanic ^e	20.9	25.4	17.1	-0.6*	-0.5	-1.0*	-	-	-
Black	14.6	17.6	12.2	-1.0*	-0.3	0.4	0.2	-0.5	-0.6
Asian/Pacific Islander	13.5	16.5	11.1	0.2	0.3	0.2	-	-	-
Amer Ind/Alaska Nat ^f	12.9	14.6	11.6	1.7*	0.8	2.4*	-	-	-
Hispanic ^e	17.7	20.7	15.3	-0.6	0.1	0.7*	-	-	-
<u>U.S. Mortality^g</u>									
RACE/ETHNICITY	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	6.0	7.7	4.7	-2.4*	-2.2*	-2.8*			
White	6.3	8.1	4.9	-2.4*	-2.1*	-2.8*			
White Hispanic ^e	5.4	6.7	4.4	-1.1*	-0.7	-1.9*			
White Non-Hispanic ^e	6.3	8.1	4.9	-2.5*	-2.5*	-2.9*			
Black	4.4	5.7	3.5	-2.3*	-2.4*	-2.4*			
Asian/Pacific Islander	4.0	4.9	3.3	-1.9*	-2.1*	-1.7*			
Amer Ind/Alaska Nat									
Total U.S.	3.2	4.1	2.5	-3.4*	-1.5	-5.2*			
CHSDA Counties	4.5	5.7	3.5	-0.6	-0.4	-4.5*			
Non-CHSDA Counties	1.7	2.2	1.3	2.2	4.4	-4.7*			
Hispanic ^e	5.1	6.3	4.2	-1.3*	-0.9	-2.1*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 19.2

SEER Incidence and US Death Rates^a Non-Hodgkin Lymphoma, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 20.19
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	11.1	16.7	6.2	0.5*	0.6*	-0.7*	0.6*	0.4	0.2
White	11.6	17.4	6.4	0.9*	0.8*	0.5	1.0*	0.8*	0.5
White Hispanic ^e	6.9	10.4	4.0	0.9	-0.6	-0.3	-	-	-
White Non-Hispanic ^e	12.5	18.8	6.9	1.1*	1.0*	0.6	-	-	-
Black	9.2	14.4	5.2	-2.5*	-2.8*	-1.7*	-2.5*	-3.0*	-1.5*
Asian/Pacific Islander	7.7	11.1	4.9	-0.9*	-0.8*	-0.8*	-	-	-
Amer Ind/Alaska Nat ^f	9.1	13.2	5.7	-1.0	-2.0	0.7	-	-	-
Hispanic ^e	6.7	10.0	4.0	0.2	-0.6	-0.4	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	2.4	3.8	1.3	-0.5	-0.3	-1.2*			
White	2.4	3.7	1.3	-0.1	0.2	-1.1*			
White Hispanic ^e	1.6	2.5	0.8	-0.5	-1.0*	-0.8*			
White Non-Hispanic ^e	2.5	3.8	1.4	0.1	0.4	-1.0*			
Black	2.9	5.0	1.3	-2.4*	-3.7*	-3.3*			
Asian/Pacific Islander	1.9	2.9	1.2	-2.1*	-2.2*	-1.6*			
Amer Ind/Alaska Nat									
Total U.S.	1.5	2.3	0.9	-2.4*	-2.7*	-2.0			
CHSDA Counties	2.1	3.3	1.1	-2.0*	-2.1*	-1.7			
Non-CHSDA Counties	0.9	1.1	0.7	-2.6	-	-			
Hispanic ^e	1.5	2.4	0.8	-0.8	-1.4*	-0.9*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 20.2

SEER Incidence and US Death Rates^a Cancer of the Oral Cavity and Pharynx, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.


Figure 20.3

SEER Incidence Rates^a

Cancer of the Oral Cavity and Pharyngeal for Selected Subsites, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints. Regression lines could not be calculated for data series that had at least one diagnosis year with a 0 case count.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Table 21.11
Cancer of the Ovary (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Females	Females	Females
All Races	11.9	-1.1*	-1.9*
White	12.5	-1.7*	-1.9*
White Hispanic ^e	11.1	-0.4	-
White Non-Hispanic ^e	12.6	-1.5*	-
Black	9.6	-0.5	-0.5*
Asian/Pacific Islander	9.3	-0.5*	-
Amer Ind/Alaska Nat ^f	10.4	-0.2	-
Hispanic ^e	10.6	-0.5	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Females	Females	
All Races	7.5	-2.2*	
White	7.8	-2.0*	
White Hispanic ^e	5.9	-0.5*	
White Non-Hispanic ^e	8.0	-2.2*	
Black	6.5	-1.1*	
Asian/Pacific Islander	4.5	-0.4	
Amer Ind/Alaska Nat			
Total U.S.	5.0	-0.4	
CHSDA Counties	6.7	0.1	
Non-CHSDA Counties	3.2	-0.6	
Hispanic ^e	5.5	-0.6*	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.


^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 21.3

SEER Incidence and US Death Rates^a Cancer of the Ovary

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473. Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 22.15
Cancer of the Pancreas (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	12.4	14.1	11.0	0.6*	0.8	0.7*	0.6*	0.7	1.0*
White	12.4	14.1	10.9	1.0*	0.8*	0.8*	0.7	1.0*	1.1*
White Hispanic ^e	11.5	12.3	10.8	0.7*	0.7*	0.6*	-	-	-
White Non-Hispanic ^e	12.5	14.4	10.9	1.0*	0.8*	0.8*	-	-	-
Black	15.5	17.2	14.2	-0.5	-0.3	-0.2	-0.1	-0.4*	0.1
Asian/Pacific Islander	10.0	11.0	9.3	0.7*	0.2	1.2*	-	-	-
Amer Ind/Alaska Nat ^f	9.7	10.9	8.6	-0.7	1.9	0.2	-	-	-
Hispanic ^e	11.1	11.8	10.4	0.6*	0.7*	0.5*	-	-	-
<u>U.S. Mortality^g</u>									
RACE/ETHNICITY	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	10.9	12.5	9.5	0.3*	0.2*	0.3*			
White	10.8	12.5	9.4	0.4*	0.4*	0.4*			
White Hispanic ^e	9.1	10.1	8.2	0.4*	0.3*	0.5*			
White Non-Hispanic ^e	11.0	12.7	9.5	0.5*	0.3*	0.4*			
Black	13.5	15.0	12.2	-0.4*	-0.6*	-0.4*			
Asian/Pacific Islander	7.7	8.3	7.3	0.1	0.3	0.5*			
Amer Ind/Alaska Nat									
Total U.S.	6.5	7.1	5.9	0.3	1.0	-0.2			
CHSDA Counties	8.7	9.6	8.0	1.5*	2.1*	1.1			
Non-CHSDA Counties	3.9	4.4	3.5	-1.7*	-0.8	-2.4*			
Hispanic ^e	8.6	9.6	7.8	0.3*	0.2	0.4*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 22.2

SEER Incidence and US Death Rates^a Cancer of the Pancreas, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 23.11
Cancer of the Prostate (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Males	Males	Males
All Races	129.4	-5.2*	-5.1*
White	121.9	-5.5*	-5.4*
White Hispanic ^e	104.6	-6.2*	-
White Non-Hispanic ^e	125.1	-5.4*	-
Black	203.5	-4.5*	-4.5*
Asian/Pacific Islander	68.9	-6.5*	-
Amer Ind/Alaska Nat ^f	63.9	-2.4*	-
Hispanic ^e	106.9	-5.7*	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Males	Males	
All Races	20.7	-3.6*	
White	19.1	-3.3*	
White Hispanic ^e	17.7	-2.5*	
White Non-Hispanic ^e	19.2	-3.4*	
Black	44.2	-4.2*	
Asian/Pacific Islander	9.1	-3.4*	
Amer Ind/Alaska Nat			
Total U.S.	15.0	-2.7*	
CHSDA Counties	19.1	-1.8*	
Non-CHSDA Counties	10.3	-3.7*	
Hispanic ^e	17.1	-2.8*	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.


^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 23.3

SEER Incidence and US Death Rates^a Cancer of the Prostate

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 24.15
Cancer of the Stomach (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	7.4	10.0	5.3	-1.2*	-1.7*	-0.7*	-1.5*	-1.8*	-0.9*
White	6.6	9.1	4.5	-1.1*	-1.6*	-0.7*	-1.0*	-1.8*	-1.0*
White Hispanic ^e	11.0	14.0	8.8	-1.4*	-1.9*	-1.0*	-	-	-
White Non-Hispanic ^e	5.8	8.3	3.7	-1.5*	-1.9*	-1.4*	-	-	-
Black	10.6	14.0	8.1	-1.7*	-2.1*	-1.1*	-1.6*	-2.1*	-0.9*
Asian/Pacific Islander	11.0	14.3	8.5	-3.0*	-3.1*	-2.7*	-	-	-
Amer Ind/Alaska Nat ^f	10.4	13.5	8.0	-0.2	-0.4	0.3	-	-	-
Hispanic ^e	10.6	13.5	8.5	-1.5*	-2.0*	-1.0*	-	-	-
<u>U.S. Mortality^g</u>									
RACE/ETHNICITY	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
	Total	Males	Females	Total	Males	Females			
All Races	3.3	4.5	2.4	-2.6*	-2.9*	-2.6*			
White	2.9	3.9	2.1	-2.5*	-2.8*	-2.7*			
White Hispanic ^e	5.7	7.5	4.4	-2.1*	-2.4*	-1.9*			
White Non-Hispanic ^e	2.6	3.5	1.8	-3.0*	-3.1*	-3.2*			
Black	6.1	8.8	4.2	-3.3*	-3.2*	-3.1*			
Asian/Pacific Islander	5.7	7.5	4.4	-3.8*	-3.7*	-3.8*			
Amer Ind/Alaska Nat									
Total U.S.	3.6	4.9	2.6	-3.0*	-2.9*	-3.3*			
CHSDA Counties	5.3	7.4	3.8	-2.1*	-2.1*	-2.2*			
Non-CHSDA Counties	1.6	2.1	1.2	-4.2*	-3.3*	-5.6*			
Hispanic ^e	5.4	7.1	4.1	-2.3*	-2.5*	-2.1*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 24.2

SEER Incidence and US Death Rates^a Cancer of the Stomach, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 25.11
Cancer of the Testis (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity

SEER Incidence	SEER 18 Areas ^b	SEER 13 Areas ^b	SEER 9 Areas ^b
	Rate 2009-2013	Trend 2004-2013 ^c	Trend 2004-2013 ^d
	Rate per 100,000 persons	AAPC (%)	AAPC (%)
RACE/ETHNICITY	Males	Males	Males
All Races	5.7	1.0*	0.8*
White	6.7	1.2*	1.0*
White Hispanic ^e	5.3	2.9*	-
White Non-Hispanic ^e	7.3	0.0	-
Black	1.5	0.7	2.0*
Asian/Pacific Islander	2.1	1.2*	-
Amer Ind/Alaska Nat ^f	4.9	2.7	-
Hispanic ^e	5.0	2.7*	-
<u>U.S. Mortality^g</u>			
	Rate 2009-2013	Trend 2004-2013 ^h	
	Rate per 100,000 persons	AAPC (%)	
RACE/ETHNICITY	Males	Males	
All Races	0.3	-0.5*	
White	0.3	-0.3	
White Hispanic ^e	0.3	1.1	
White Non-Hispanic ^e	0.3	-0.7*	
Black	0.1	-1.0	
Asian/Pacific Islander	0.1	0.2	
Amer Ind/Alaska Nat			
Total U.S.	0.2	-	
CHSDA Counties	0.3	-	
Non-CHSDA Counties	-	-	
Hispanic ^e	0.3	0.9	

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.


^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

* The APC is significantly different from zero (p<.05).

Figure 25.3

SEER Incidence and US Death Rates^a Cancer of the Testis

Joinpoint Analyses for Whites and Blacks from 1975-2013
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

^a Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Joinpoint could not fit a regression line for American Indian/Alaska Native Mortality due to insufficient data.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 26.15
Cancer of the Thyroid (Invasive)

SEER Incidence and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	13.9	6.9	20.6	4.4*	5.0*	4.5*	4.5*	5.3*	4.4*
White	14.7	7.4	21.9	4.4*	5.1*	4.4*	4.4*	4.5*	4.4*
White Hispanic ^e	12.4	5.2	19.5	4.9*	4.1*	5.2*	-	-	-
White Non-Hispanic ^e	15.6	8.1	23.0	4.4*	4.5*	4.4*	-	-	-
Black	8.3	3.6	12.4	4.5*	3.8*	5.0*	5.9*	3.1*	5.9*
Asian/Pacific Islander	13.6	6.5	19.9	4.7*	5.9*	4.3*	-	-	-
Amer Ind/Alaska Nat ^f	8.8	3.6	13.6	3.8*	2.3	4.1*	-	-	-
Hispanic ^e	11.9	5.0	18.6	4.8*	4.7*	5.0*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	0.5	0.5	0.5	0.8*	1.5*	0.4*			
White	0.5	0.5	0.5	0.7*	1.5*	0.3			
White Hispanic ^e	0.6	0.5	0.7	0.1	1.2	-0.5			
White Non-Hispanic ^e	0.5	0.5	0.5	0.7*	1.5*	0.3			
Black	0.5	0.4	0.6	0.9*	1.4*	0.8			
Asian/Pacific Islander	0.7	0.6	0.8	-0.2	0.7	0.0			
Amer Ind/Alaska Nat									
Total U.S.	0.4	0.4	0.4	2.1	-	-0.5			
CHSDA Counties	0.5	-	0.5	2.1	-	-0.4			
Non-CHSDA Counties	-	-	-	-	-	-			
Hispanic ^e	0.6	0.5	0.6	0.0	1.1	-0.7			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

^c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

^d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^g US Mortality Files, National Center for Health Statistics, CDC.

^h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 26.2

SEER Incidence and US Death Rates^a Cancer of the Thyroid, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Table 27.15
Cancer of the Urinary Bladder

SEER Incidence (Invasive and *In Situ*) and U.S. Mortality
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b			SEER 13 Areas ^b			SEER 9 Areas ^b		
	Rate 2009-2013			Trend 2004-2013 ^c			Trend 2004-2013 ^d		
	Rate per 100,000 persons			AAPC (%)			AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	20.1	35.3	8.6	-1.0*	-1.1*	-1.3*	-0.8*	-0.9*	-1.2*
White	22.0	38.5	9.2	-0.9*	-1.1*	-1.3*	-0.7*	-0.9*	-1.0*
White Hispanic ^e	11.4	19.9	5.2	-0.3	-0.3	-0.6	-	-	-
White Non-Hispanic ^e	23.6	41.3	9.9	-0.7*	-0.9*	-1.1*	-	-	-
Black	12.6	21.2	6.9	0.2	0.5*	-0.4	0.2*	0.4*	0.1
Asian/Pacific Islander	8.8	15.5	3.9	-1.7*	-1.8*	-0.2	-	-	-
Amer Ind/Alaska Nat ^f	8.1	14.4	3.5	1.2	1.4	3.6	-	-	-
Hispanic ^e	11.1	19.4	5.0	-0.4	-0.3	-0.6*	-	-	-
<u>U.S. Mortality^g</u>									
	Rate 2009-2013			Trend 2004-2013 ^h					
	Rate per 100,000 persons			AAPC (%)					
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females			
All Races	4.4	7.7	2.2	0.0	0.0	-0.5*			
White	4.6	8.1	2.2	0.3*	0.2*	-0.3*			
White Hispanic ^e	2.5	4.2	1.3	0.0	-0.2	-0.1			
White Non-Hispanic ^e	4.8	8.4	2.3	0.3*	0.3*	-0.2*			
Black	3.6	5.4	2.5	-0.9*	-0.8*	-1.3*			
Asian/Pacific Islander	1.7	2.9	0.9	-0.6	0.0	-1.3*			
Amer Ind/Alaska Nat									
Total U.S.	1.8	3.1	1.0	1.1	1.5	-0.3			
CHSDA Counties	2.1	3.8	1.1	2.4*	3.5*	-			
Non-CHSDA Counties	1.5	2.4	0.9	-1.0	-1.1	-			
Hispanic ^e	2.3	4.0	1.2	-0.2	-0.3	-0.2			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

c The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

d The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

f Incidence data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

g US Mortality Files, National Center for Health Statistics, CDC.

h The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.


* The APC is significantly different from zero (p<.05).

Figure 27.2

SEER Incidence and US Death Rates^a Cancer of the Urinary Bladder, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.


- ^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.
- ^b API = Asian/Pacific Islander.
- ^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Figure 28.3

SEER Incidence and US Death Rates^a All Cancer Sites, Ages <20, Both Sexes

Joinpoint Analyses for Whites and Blacks from 1975-2013

and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2013



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from US Mortality Files, National Center for Health Statistics, CDC.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. Joinpoint analyses for Whites and Blacks during the 1975-2013 period allow a maximum of 5 joinpoints. Analyses for other ethnic groups during the period 1992-2013 allow a maximum of 4 joinpoints.


^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from New Hampshire and Oklahoma.

Figure 29.2

Childhood Cancer: SEER Incidence Rates 2009- 2013 by ICCC Group and Race/Ethnicity (includes myelodysplastic syndromes and Group III benign brain) Both Sexes, Under 20 Years of Age



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Incidence rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.


^a Rate for Group III (Brain/CNS) includes benign brain tumors.

^b Incidence rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Figure 29.3

Childhood Cancer 5-Year Observed Survival (%) by ICCC Group^a and Race/Ethnicity Both Sexes, Under 20 Years of Age, 2006-2012



Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005; Vol 103, No. 7, pg 1457-1467.

^a Excludes myelodysplastic syndromes (MDS) and benign brain tumors.

^b American Indian/Alaska Native. Survival data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Survival data for Hispanics are based on NHIA and excludes cases from the Alaska Native Registry.