

Trends in SEER 9 Observed Incidence^a Using the Joinpoint Regression Program, 1975-2013 With up to Five Joinpoints,

By Race and Sex

| | | JP Trend 1 | | JP Trend 2 | | JP Trend 3 | | JP Trend 4 | | JP Trend 5 | | JP Trend 6 | | _AAPC ^d _ |
|---|------------|------------|------|------------|-------|------------|---------------|------------|-------|------------|-------|------------|-------|----------------------|
| - | | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | Years | APC | 2009-13 |
| | | | | | | | | | | | | | | |
| SEER 9 Observed Incidence ^a for Cancer of the Colon, 1975-2013 | | | | | | | | | | | | | | |
| All Races | Both Sexes | 1975-85 | 1.1* | 1985-95 | -1.6* | 1995-01 | -0.1 | 2001-13 | -3.3* | | | | | -3.3* |
| All Races | Male | 1975-85 | 1.7* | 1985-95 | -1.0" | 1991-95 | -0.1 -3.0* | 1995-98 | 1.3 | 1998-04 | -2.5* | 2004-13 | -3.7* | -3.7* |
| | | | | | | | | | | 1998-04 | -2.5^ | 2004-13 | -3./~ | |
| All Races | Female | 1975-85 | 0.6* | 1985-93 | -1.8* | 1993-02 | -0.3 | 2002-13 | -3.2* | | | | | -3.2* |
| White | Both Sexes | 1975-85 | 1.1* | 1985-95 | -1.9* | 1995-98 | 1.3 | 1998-07 | -2.4* | 2007-13 | -4.2* | | | -4.2* |
| White | Male | 1975-85 | 1.7* | 1985-91 | -1.4* | 1991-95 | -3.1* | 1995-98 | 1.6 | 1998-08 | -3.0* | 2008-13 | -4.7* | -4.7* |
| White | Female | 1975-85 | 0.6* | 1985-93 | -2.1* | 1993-01 | -0.1 | 2001-13 | -3.1* | 1000 00 | 3.0 | 2000 13 | 1., | -3.1* |
| WIIICE | remare | 1973-03 | 0.0 | 1905-93 | -2.1 | 1993-01 | -0.1 | 2001-13 | -3.1 | | | | | -3.1 |
| Black | Both Sexes | 1975-80 | 3.6* | 1980-03 | -0.1 | 2003-13 | -3.6* | | | | | | | -3.6* |
| Black | Male | 1975-87 | 1.8* | 1987-03 | -0.4 | 2003-13 | -3.4* | | | | | | | -3.4* |
| Black | Female | 1975-04 | -0.1 | 2004-13 | -4.3* | | | | | | | | | -4.3* |
| | | | | | | | | | | | | | | |
| SEER 9 Observed Incidence for Cancer of the Rectum, 1975-2013 | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | |
| All Races | Both Sexes | 1975-87 | -0.3 | 1987-95 | -2.4* | 1995-98 | 2.7 | 1998-13 | -2.3* | | | | | -2.3* |
| All Races | Male | 1975-89 | -0.3 | 1989-94 | -3.6* | 1994-98 | 1.7 | 1998-13 | -2.6* | | | | | -2.6* |
| All Races | Female | 1975-84 | -0.2 | 1984-95 | -2.2* | 1995-98 | 2.6 | 1998-13 | -2.2* | | | | | -2.2* |
| | | | | | | | | | | | | | | |
| White | Both Sexes | 1975-85 | -0.1 | 1985-95 | -2.3* | 1995-98 | 2.5 | 1998-13 | -2.7* | | | | | -2.7* |
| White | Male | 1975-89 | -0.3 | 1989-94 | -3.9* | 1994-99 | 1.1 | 1999-13 | -3.1* | | | | | -3.1* |
| White | Female | 1975-84 | -0.2 | 1984-95 | -2.3* | 1995-98 | 2.5 | 1998-13 | -2.5* | | | | | -2.5* |
| ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, | 10 | 2373 01 | 0.2 | 1701 73 | 2.5 | 1333 33 | 2.5 | 1330 13 | 2.5 | | | | | 2.0 |
| Black | Both Sexes | 1975-08 | 0.0 | 2008-13 | -4.4* | | | | | | | | | -4.4* |
| Black | Male | 1975-08 | 0.0 | 2008-13 | -3.8 | | | | | | | | | -3.8 |
| Black | Female | 1975-09 | -0.1 | 2009-13 | -7.4* | | | | | | | | | -7.4* |
| | | | | | | | | | | | | | | |

Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute. (http://surveillance.cancer.gov/joinpoint/). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^{*} The APC/AAPC is significantly different from zero (p<.05).

⁻ Joinpoint regression line analysis could not be performed on data series.