

Table 6.5  
Cancer of the Colon and Rectum (Invasive)

Delay-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	59.54	68.45	53.66	60.19	69.76	54.05	56.85	59.33	54.66
1976	61.34	71.72	54.12	62.14	72.68	55.00	55.04	58.53	51.28
1977	62.40	73.48	54.95	63.19	74.32	55.75	60.69	75.66	51.47
1978	62.03	72.34	55.25	62.77	73.87	55.59	62.10	65.83	60.42
1979	62.37	74.11	54.58	62.94	75.19	54.94	58.57	61.79	56.16
1980	63.76	74.56	56.54	63.95	74.88	56.83	70.83	79.71	64.23
1981	64.25	75.82	56.49	64.87	76.68	57.14	62.83	72.89	55.97
1982	62.79	75.18	54.51	63.44	76.24	55.05	61.25	68.97	55.92
1983	63.65	75.54	55.61	64.24	76.51	56.04	66.79	75.80	61.13
1984	64.82	77.65	56.08	65.83	79.26	56.74	62.46	70.42	58.20
1985	66.30	79.19	57.31	67.19	80.20	58.35	64.06	74.39	56.39
1986	64.18	77.78	54.70	64.57	78.88	54.72	64.66	73.09	58.75
1987	62.71	77.47	52.68	62.89	78.18	52.63	66.28	77.63	59.01
1988	61.37	75.19	51.80	61.54	75.77	51.73	63.00	72.41	56.62
1989	61.70	74.98	52.42	61.94	75.42	52.64	64.52	79.41	54.76
1990	60.70	74.36	51.28	60.66	74.75	51.03	66.87	75.15	61.66
1991	59.48	72.91	50.04	59.29	73.11	49.65	65.65	76.42	58.16
1992	58.00	70.41	49.21	57.81	70.66	48.68	65.66	78.59	57.57
1993	56.79	68.43	48.30	56.31	68.06	47.79	64.49	78.12	55.25
1994	55.63	66.28	47.73	55.37	66.32	47.17	64.44	75.27	57.98
1995	54.04	63.84	46.91	53.68	63.40	46.58	62.04	71.49	55.98
1996	54.78	65.60	46.67	54.73	65.94	46.34	58.63	66.66	53.11
1997	56.40	68.08	47.92	56.35	67.87	47.99	64.58	77.55	56.17
1998	56.78	67.07	49.31	56.66	67.19	48.99	63.31	74.49	56.30
1999	55.48	65.75	47.84	55.28	65.85	47.18	63.21	71.89	58.15
2000	54.12	63.65	46.76	53.73	63.02	46.46	64.37	74.04	58.06
2001	53.59	63.05	46.20	52.95	62.60	45.31	64.35	74.94	57.59
2002	53.17	61.54	46.64	52.27	60.28	45.88	63.22	74.04	56.59
2003	50.80	59.49	43.93	50.22	58.55	43.46	62.57	75.31	54.68
2004	49.88	58.60	43.16	49.04	57.87	42.08	62.43	74.43	54.53
2005	47.85	55.30	41.91	47.41	55.10	41.16	58.39	65.14	54.71
2006	46.91	53.34	41.70	46.16	52.65	40.73	58.24	65.33	53.64
2007	46.48	53.42	40.79	45.57	52.25	40.03	57.94	65.83	52.30
2008	45.38	51.66	40.17	44.57	50.56	39.54	55.17	65.04	48.00
2009	43.38	49.07	38.71	42.08	47.58	37.43	54.17	61.40	49.66
2010	41.09	47.22	36.11	39.80	45.88	34.71	49.99	57.26	45.23
2011	39.87	45.70	35.09	38.77	43.99	34.37	48.59	56.49	43.07
2012	39.05	44.25	34.65	38.19	42.81	34.15	48.29	57.40	41.85
2013	37.87	43.72	32.97	36.71	41.54	32.61	47.77	59.17	39.68

Additional information on the model used to delay-adjust SEER Incidence rates can be found at <http://surveillance.cancer.gov/delay/>.

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.