

Table 6.29

Cancer of the Colon And Rectum (Invasive)

Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2009-2013
Both Sexes by Race

| Histology ^a | All Races | | White | | Black | | Asian/Pacific Islander | | American Indian/ Alaska Native ^b | | Hispanic ^c | |
|--|-----------|---------|---------|---------|--------|---------|---------------------------|---------|--|---------|-----------------------|---------|
| | Count | Percent | Count | Percent | Count | Percent | Count | Percent | Count | Percent | Count | Percent |
| Carcinoma | 178,971 | 99.4% | 139,391 | 99.4% | 21,893 | 99.4% | 14,568 | 99.4% | 1,018 | 99.6% | 18,902 | 99.4% |
| Epidermoid carcinoma ^d (8051-8131) | 1,122 | 0.6% | 958 | 0.7% | 116 | 0.5% | 25 | 0.2% | - | - | 97 | 0.5% |
| Squamous cell carcinoma (8070-8078,8083-8084) | 1,069 | 0.6% | 918 | 0.7% | 110 | 0.5% | 24 | 0.2% | - | - | 94 | 0.5% |
| Adenocarcinoma (8050,8140-8147, 8160-8162,8180-8221, 8250-8507,8520-8551,8560, 8570-8574,8576,8940-8941) | 167,109 | 92.8% | 131,344 | 93.7% | 19,763 | 89.8% | 13,494 | 92.1% | 946 | 92.6% | 17,541 | 92.2% |
| Adenocarcinoma, NOS (8140) | 114,952 | 63.8% | 90,084 | 64.3% | 13,712 | 62.3% | 9,625 | 65.7% | 675 | 66.0% | 12,150 | 63.9% |
| Adenocarcinoma in adenomatous polyp (8210) | 15,085 | 8.4% | 11,699 | 8.3% | 1,701 | 7.7% | 1,309 | 8.9% | 105 | 10.3% | 1,343 | 7.1% |
| Adenocarcinoma in villous adenoma (8261) | 3,977 | 2.2% | 3,184 | 2.3% | 440 | 2.0% | 299 | 2.0% | 16 | 1.6% | 535 | 2.8% |
| Mucinous adenocarcinoma (8470-8473,8480,8482) | 11,517 | 6.4% | 9,477 | 6.8% | 1,192 | 5.4% | 737 | 5.0% | 50 | 4.9% | 1,335 | 7.0% |
| Mucin-producing adenocarcinoma (8481) | 1,571 | 0.9% | 1,251 | 0.9% | 211 | 1.0% | 100 | 0.7% | - | - | 155 | 0.8% |
| Signet ring adenocarcinoma (8490) | 2,082 | 1.2% | 1,707 | 1.2% | 198 | 0.9% | 152 | 1.0% | - | - | 227 | 1.2% |
| Other adenocarcinomas ^e | 17,925 | 10.0% | 13,942 | 9.9% | 2,309 | 10.5% | 1,272 | 8.7% | 90 | 8.8% | 1,796 | 9.4% |
| Other specific carcinomas (8012-8015,8030-8046, 8150-8155,8170-8175, 8230-8249,8508,8510-8513, 8561-8562,8575,8580-8671) | 9,207 | 5.1% | 5,901 | 4.2% | 1,813 | 8.2% | 946 | 6.5% | 53 | 5.2% | 1,090 | 5.7% |
| Neuroendocrine carcinoid (8240-8245) | 6,785 | 3.8% | 4,111 | 2.9% | 1,443 | 6.6% | 762 | 5.2% | 44 | 4.3% | 858 | 4.5% |
| Other ^f | 2,422 | 1.3% | 1,790 | 1.3% | 370 | 1.7% | 184 | 1.3% | - | - | 232 | 1.2% |
| Unspecified, Carcinoma, NOS (8010-8011, 8020-8022) | 1,533 | 0.9% | 1,188 | 0.8% | 201 | 0.9% | 103 | 0.7% | 16 | 1.6% | 174 | 0.9% |

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Percents may not sum to 100 due to rounding.

^a Excludes Kaposi Sarcoma, mesothelioma, lymphomas, leukemias, myelomas, lymphoreticular, and immunoproliferative diseases.

^b Estimates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^d Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^e Epidermoid carcinoma includes squamous, basal, and transitional cell carcinomas.

^f Includes histologies 8050, 8141-8147, 8160-8162, 8180-8204, 8211-8221, 8250-8260, 8262-8264, 8270-8463, 8500-8507, 8514, 8520-8551, 8560, 8570-8574, 8576, 8940-8941.

^f Includes histologies 8012-8015, 8030-8046, 8150-8155, 8170-8175, 8230-8231, 8246-8249, 8508, 8510-8513, 8561-8562, 8575, 8580-8671.

- Statistic not shown due to fewer than 16 cases during the time period.

Table 6.29 - continued

Cancer of the Colon And Rectum (Invasive)

Percent Distribution and Counts by Histology among Histologically Confirmed Cases, 2009-2013
Both Sexes by Race

| Histology ^a | All Races | | White | | Black | | Asian/Pacific Islander | | American Indian/ Alaska Native ^b | | Hispanic ^c | |
|---|-----------|---------|---------|---------|--------|---------|------------------------|---------|--|---------|-----------------------|---------|
| | Count | Percent | Count | Percent | Count | Percent | Count | Percent | Count | Percent | Count | Percent |
| Sarcoma and other soft tissue tumors (8680-8713,8800-8921, 8990-8991,9040-9044, 9120-9136,9150-9252, 9370-9373,9540-9582) | 108 | 0.1% | 88 | 0.1% | - | - | - | - | - | - | 21 | 0.1% |
| Other specific types (8720-8790,8930-8936, 8950-8983,9000-9030, 9060-9110,9260-9365, 9380-9539) | 293 | 0.2% | 200 | 0.1% | 49 | 0.2% | 43 | 0.3% | - | - | 33 | 0.2% |
| Unspecified (8000-8005) | 670 | 0.4% | 496 | 0.4% | 65 | 0.3% | 41 | 0.3% | - | - | 69 | 0.4% |
| Total ^d | 180,042 | 100.0% | 140,175 | 100.0% | 22,019 | 100.0% | 14,658 | 100.0% | 1,022 | 100.0% | 19,025 | 100.0% |

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Percents may not sum to 100 due to rounding.

^a Excludes Kaposi Sarcoma, mesothelioma, lymphomas, leukemias, myelomas, lymphoreticular, and immunoproliferative diseases.

^b Estimates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Total includes carcinoma cases from the previous page.

- Statistic not shown due to fewer than 16 cases during the time period.