

Table 7.4  
Cancer of the Corpus and Uterus, NOS (Invasive)

Delay-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Age

|                    | All Races, Females |      |        | White Females |      |        | Black Females |      |       |
|--------------------|--------------------|------|--------|---------------|------|--------|---------------|------|-------|
|                    | All                | <50  | 50+    | All           | <50  | 50+    | All           | <50  | 50+   |
| Year of Diagnosis: |                    |      |        |               |      |        |               |      |       |
| 1975               | 35.47              | 6.48 | 111.37 | 37.32         | 6.81 | 117.22 | 19.91         | 3.64 | 62.51 |
| 1976               | 34.37              | 6.20 | 108.12 | 36.26         | 6.28 | 114.75 | 17.61         | 3.29 | 55.11 |
| 1977               | 31.67              | 5.75 | 99.55  | 33.18         | 5.93 | 104.52 | 20.06         | 3.52 | 63.37 |
| 1978               | 30.05              | 6.10 | 92.76  | 31.66         | 6.31 | 98.05  | 18.74         | 3.94 | 57.49 |
| 1979               | 27.91              | 5.33 | 87.03  | 29.24         | 5.35 | 91.82  | 16.96         | 3.94 | 51.06 |
| 1980               | 27.31              | 4.97 | 85.81  | 28.53         | 4.95 | 90.28  | 16.22         | 2.51 | 52.13 |
| 1981               | 27.27              | 5.60 | 83.99  | 28.35         | 5.45 | 88.32  | 17.42         | 2.86 | 55.52 |
| 1982               | 26.80              | 5.06 | 83.74  | 28.23         | 5.22 | 88.49  | 17.09         | 2.49 | 55.33 |
| 1983               | 26.51              | 4.86 | 83.21  | 27.77         | 4.89 | 87.66  | 18.83         | 3.39 | 59.25 |
| 1984               | 25.70              | 4.62 | 80.89  | 27.15         | 4.89 | 85.44  | 17.38         | 1.74 | 58.35 |
| 1985               | 25.23              | 4.63 | 79.19  | 26.56         | 4.83 | 83.45  | 17.41         | 2.23 | 57.17 |
| 1986               | 24.38              | 4.81 | 75.64  | 25.61         | 4.97 | 79.66  | 16.78         | 2.54 | 54.09 |
| 1987               | 24.71              | 5.00 | 76.32  | 26.12         | 5.28 | 80.68  | 15.76         | 2.02 | 51.73 |
| 1988               | 23.58              | 4.56 | 73.38  | 24.62         | 4.49 | 77.34  | 16.97         | 2.97 | 53.65 |
| 1989               | 24.43              | 4.48 | 76.68  | 25.50         | 4.77 | 79.80  | 19.36         | 1.78 | 65.40 |
| 1990               | 25.05              | 4.60 | 78.60  | 26.55         | 4.82 | 83.44  | 17.03         | 2.28 | 55.65 |
| 1991               | 24.63              | 4.37 | 77.70  | 25.99         | 4.48 | 82.33  | 17.48         | 3.12 | 55.09 |
| 1992               | 24.78              | 4.86 | 76.96  | 26.15         | 4.85 | 81.90  | 17.03         | 3.01 | 53.74 |
| 1993               | 24.30              | 5.19 | 74.32  | 25.68         | 5.14 | 79.48  | 16.97         | 3.48 | 52.27 |
| 1994               | 25.12              | 5.02 | 77.74  | 26.32         | 5.06 | 81.99  | 17.65         | 2.42 | 57.53 |
| 1995               | 25.32              | 4.72 | 79.27  | 26.50         | 4.64 | 83.75  | 18.82         | 2.75 | 60.90 |
| 1996               | 24.93              | 4.52 | 78.38  | 25.99         | 4.39 | 82.54  | 19.42         | 3.08 | 62.21 |
| 1997               | 25.88              | 5.04 | 80.43  | 27.33         | 5.04 | 85.70  | 16.63         | 2.54 | 53.51 |
| 1998               | 25.73              | 4.83 | 80.45  | 27.02         | 4.77 | 85.29  | 19.51         | 3.15 | 62.36 |
| 1999               | 25.32              | 5.19 | 78.05  | 26.77         | 5.06 | 83.60  | 17.79         | 2.97 | 56.60 |
| 2000               | 24.82              | 5.23 | 76.13  | 26.21         | 5.37 | 80.80  | 18.02         | 3.00 | 57.35 |
| 2001               | 25.29              | 5.11 | 78.15  | 26.69         | 4.95 | 83.62  | 19.44         | 4.07 | 59.70 |
| 2002               | 24.93              | 5.07 | 76.95  | 25.30         | 4.59 | 79.56  | 23.41         | 4.87 | 71.96 |
| 2003               | 24.37              | 5.32 | 74.25  | 25.58         | 5.31 | 78.67  | 20.26         | 3.76 | 63.46 |
| 2004               | 25.11              | 5.57 | 76.27  | 26.05         | 5.28 | 80.46  | 20.65         | 3.85 | 64.64 |
| 2005               | 24.98              | 5.54 | 75.89  | 25.86         | 5.39 | 79.46  | 23.23         | 4.12 | 73.28 |
| 2006               | 24.62              | 5.14 | 75.64  | 26.04         | 5.04 | 81.03  | 19.37         | 3.71 | 60.39 |
| 2007               | 25.34              | 5.88 | 76.29  | 26.15         | 5.90 | 79.18  | 22.73         | 3.52 | 73.02 |
| 2008               | 26.15              | 6.02 | 78.84  | 26.72         | 5.76 | 81.59  | 24.41         | 4.32 | 77.02 |
| 2009               | 27.45              | 6.02 | 83.57  | 28.09         | 5.95 | 86.08  | 25.48         | 3.74 | 82.39 |
| 2010               | 27.54              | 6.31 | 83.12  | 28.19         | 6.11 | 86.01  | 24.96         | 3.91 | 80.11 |
| 2011               | 27.24              | 5.60 | 83.92  | 27.46         | 5.38 | 85.28  | 28.56         | 4.10 | 92.61 |
| 2012               | 27.67              | 6.24 | 83.76  | 28.42         | 6.03 | 87.03  | 25.80         | 4.39 | 81.86 |
| 2013               | 27.15              | 5.95 | 82.68  | 27.27         | 5.36 | 84.65  | 27.85         | 5.00 | 87.70 |

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.