

Table 7.15  
Cancer of the Corpus and Uterus, NOS (Invasive)

Estimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2013  
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 <sup>e</sup>	0 to <38 <sup>e</sup>	>=38 <sup>g</sup>	Complete <sup>h</sup>
<u>Race</u>										
All Races <sup>b</sup>	190,409	131,077	102,712	73,340	48,841	32,618	529,671	615,772	19,665	635,437
White <sup>b</sup>	161,701	116,009	92,657	66,999	45,395	30,608	467,264	548,026	16,348	564,374
Black <sup>b</sup>	17,049	8,164	5,300	3,088	1,771	1,058	34,703	37,414	672	38,086
Asian/Pacific Islander <sup>c</sup>	7,713	4,660	3,176	+	+	+	17,931	+	+	+
Hispanic <sup>d</sup>	15,050	8,909	5,628	+	+	+	33,874	+	+	+

Estimated prevalence percent<sup>a</sup> on January 1, 2013, of the SEER<sup>c</sup> population diagnosed in the previous 21 years  
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted <sup>f</sup>
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races <sup>c</sup>	0.2867%	-	-	0.0034%	0.0314%	0.1217%	0.3988%	0.9484%	1.2559%	1.0819%	0.2471%
White <sup>c</sup>	0.3261%	-	-	0.0033%	0.0314%	0.1211%	0.4156%	1.0303%	1.4062%	1.2172%	0.2681%
Black <sup>c</sup>	0.1507%	-	-	0.0036%	0.0236%	0.0882%	0.2262%	0.6325%	0.8881%	0.5960%	0.1618%
Asian/Pacific Islander <sup>c</sup>	0.2089%	-	-	0.0034%	0.0316%	0.1414%	0.4355%	0.7171%	0.6716%	0.5085%	0.1835%
Hispanic <sup>d</sup>	0.1324%	-	-	0.0047%	0.0418%	0.1318%	0.3325%	0.6877%	0.8177%	0.6328%	0.1841%

<sup>a</sup> US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

<sup>e</sup> Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <38 and >=38. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.