

Table 14.13
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.01	0.01	1.35	0.97	Asian/ Pacific Islander	0	0.01	0.01	0.01	2.56	2.06
	10	0.00	0.00	0.01	1.36	0.98		10	0.00	0.01	0.03	2.58	2.08
	20	0.00	0.01	0.07	1.36	0.99		20	0.00	0.03	0.15	2.58	2.08
	30	0.01	0.07	0.37	1.38	1.00		30	0.02	0.14	0.52	2.59	2.09
	40	0.06	0.37	0.80	1.39	1.01		40	0.12	0.50	1.11	2.59	2.09
	50	0.32	0.76	1.12	1.37	1.01		50	0.38	1.00	1.73	2.51	2.05
	60	0.48	0.87	1.09	1.14	0.89		60	0.65	1.40	2.01	2.21	1.87
	70	0.45	0.72	-	0.77	0.68		70	0.83	1.49	-	1.71	1.60
80	0.38	-	-	0.45	0.46	80	0.82	-	-	1.09	1.18		
White	0	0.01	0.01	0.01	1.21	0.90	American Indian/ Alaska Native ^a	0	0.02	0.02	0.02	1.67	1.35
	10	0.00	0.00	0.01	1.22	0.91		10	0.00	0.00	0.01	1.67	1.36
	20	0.00	0.01	0.06	1.22	0.92		20	0.00	0.01	0.10	1.68	1.38
	30	0.01	0.06	0.33	1.23	0.93		30	0.01	0.10	0.52	1.72	1.41
	40	0.05	0.33	0.71	1.25	0.94		40	0.10	0.52	1.09	1.76	1.45
	50	0.29	0.68	1.01	1.23	0.94		50	0.45	1.04	1.55	1.74	1.49
	60	0.41	0.77	0.98	1.01	0.83		60	0.65	1.20	1.42	1.42	1.37
	70	0.42	0.66	-	0.70	0.65		70	0.65	0.90	-	0.90	1.04
80	0.34	-	-	0.40	0.44	80	0.36	-	-	0.37	0.65		
Black	0	0.01	0.01	0.01	1.48	1.18	Hispanic ^b	0	0.01	0.01	0.01	2.25	1.63
	10	0.00	0.01	0.02	1.51	1.21		10	0.00	0.00	0.01	2.27	1.65
	20	0.00	0.02	0.07	1.52	1.22		20	0.00	0.01	0.08	2.27	1.66
	30	0.01	0.06	0.47	1.54	1.24		30	0.01	0.08	0.50	2.29	1.67
	40	0.05	0.47	1.14	1.56	1.26		40	0.08	0.50	1.15	2.31	1.69
	50	0.44	1.14	1.42	1.58	1.28		50	0.44	1.10	1.73	2.29	1.68
	60	0.78	1.10	1.25	1.28	1.10		60	0.70	1.37	1.81	1.96	1.53
	70	0.40	0.60	-	0.63	0.70		70	0.76	1.26	-	1.43	1.22
80	0.31	-	-	0.36	0.48	80	0.66	-	-	0.89	0.86		

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.