



Table 14.18
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	6.14	0.02		
High Five States				
D.C.	9.59	0.57	01(01,02)	56.31 ^c
Louisiana	7.86	0.18	02(02,06)	28.04 ^c
Texas	7.81	0.08	03(02,04)	27.25 ^c
Hawaii	7.70	0.31	04(01,08)	25.45 ^c
California	7.43	0.06	05(03,07)	21.11 ^c
Low Five States				
Montana	4.69	0.28	47(32,51)	-23.49 ^c
Idaho	4.60	0.23	48(36,51)	-25.07 ^c
South Dakota	4.58	0.31	49(32,51)	-25.34 ^c
Utah	4.55	0.20	50(38,51)	-25.82 ^c
North Dakota	3.76	0.31	51(46,51)	-38.80 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	6.34	0.15	14(07,22)	3.36	Montana	4.69	0.28	47(32,51)	-23.49 ^c
Alaska	5.92	0.47	21(05,48)	-3.50	Nebraska	4.81	0.22	45(33,50)	-21.58 ^c
Arizona	6.00	0.13	20(13,30)	-2.23	Nevada	5.66	0.20	29(14,41)	-7.82
Arkansas	6.56	0.20	11(06,20)	6.99	New Hampshire	5.00	0.25	42(25,50)	-18.46 ^c
California	7.43	0.06	05(03,07)	21.11 ^c	New Jersey	5.63	0.11	30(20,38)	-8.20
Colorado	5.27	0.15	37(26,46)	-14.07	New Mexico	6.86	0.25	07(04,17)	11.77
Connecticut	5.43	0.16	34(22,43)	-11.45	New York	6.05	0.07	18(14,25)	-1.41
Delaware	6.61	0.35	09(04,30)	7.67	North Carolina	5.71	0.10	27(19,36)	-7.01
D.C.	9.59	0.57	01(01,02)	56.31 ^c	North Dakota	3.76	0.31	51(46,51)	-38.80 ^c
Florida	5.84	0.07	24(18,31)	-4.82	Ohio	5.47	0.09	33(25,40)	-10.88
Georgia	5.75	0.11	26(17,35)	-6.25	Oklahoma	6.01	0.17	19(10,33)	-2.04
Hawaii	7.70	0.31	04(01,08)	25.45 ^c	Oregon	6.34	0.17	15(07,24)	3.29
Idaho	4.60	0.23	48(36,51)	-25.07 ^c	Pennsylvania	5.68	0.08	28(20,36)	-7.38
Illinois	5.86	0.09	22(16,31)	-4.41	Rhode Island	6.67	0.33	08(04,28)	8.67
Indiana	5.38	0.12	35(26,43)	-12.26	South Carolina	5.86	0.15	23(14,35)	-4.52
Iowa	4.75	0.16	46(37,50)	-22.55 ^c	South Dakota	4.58	0.31	49(32,51)	-25.34 ^c
Kansas	5.10	0.18	41(29,48)	-16.83 ^c	Tennessee	6.42	0.13	13(07,20)	4.59
Kentucky	5.82	0.15	25(14,36)	-5.09	Texas	7.81	0.08	03(02,04)	27.25 ^c
Louisiana	7.86	0.18	02(02,06)	28.04 ^c	Utah	4.55	0.20	50(38,51)	-25.82 ^c
Maine	5.12	0.24	39(22,50)	-16.54 ^c	Vermont	5.16	0.37	38(14,50)	-15.88 ^c
Maryland	6.25	0.14	16(08,24)	1.85	Virginia	5.60	0.11	31(20,38)	-8.67
Massachusetts	6.46	0.13	12(07,18)	5.32	Washington	6.57	0.14	10(06,17)	7.12
Michigan	5.55	0.10	32(23,38)	-9.51	West Virginia	4.92	0.20	44(32,50)	-19.85 ^c
Minnesota	5.00	0.13	43(35,48)	-18.47 ^c	Wisconsin	5.29	0.13	36(27,44)	-13.71
Mississippi	7.12	0.21	06(03,11)	15.98 ^c	Wyoming	5.12	0.41	40(12,51)	-16.60 ^c
Missouri	6.07	0.13	17(11,28)	-1.02					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.