


Table 14.19
 Cancer of the Liver and Intrahepatic Bile Duct (Invasive)
 Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	9.07	0.03		
High Five States				
D.C.	14.97	1.08	01(01,02)	65.04 ^c
Louisiana	11.81	0.33	02(01,06)	30.21 ^c
Texas	11.54	0.15	03(02,05)	27.24 ^c
Hawaii	11.31	0.55	04(02,12)	24.73 ^c
California	10.87	0.11	05(04,08)	19.83 ^c
Low Five States				
Idaho	6.46	0.41	47(36,51)	-28.73 ^c
Montana	6.38	0.47	48(35,51)	-29.64 ^c
Utah	5.97	0.34	49(44,51)	-34.18 ^c
South Dakota	5.96	0.52	50(38,51)	-34.29 ^c
North Dakota	5.40	0.55	51(42,51)	-40.42 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	9.39	0.28	14(07,26)	3.54	Montana	6.38	0.47	48(35,51)	-29.64 ^c
Alaska	8.51	0.82	25(04,49)	-6.14	Nebraska	7.08	0.39	44(29,50)	-21.92 ^c
Arizona	8.41	0.22	30(17,37)	-7.31	Nevada	7.94	0.35	34(18,46)	-12.42
Arkansas	9.79	0.36	11(05,23)	7.93	New Hampshire	7.74	0.47	37(17,48)	-14.66
California	10.87	0.11	05(04,08)	19.83 ^c	New Jersey	8.49	0.20	28(17,36)	-6.44
Colorado	7.78	0.26	36(25,44)	-14.23	New Mexico	9.94	0.43	10(04,23)	9.57
Connecticut	8.50	0.30	26(14,39)	-6.34	New York	9.09	0.14	18(12,24)	0.18
Delaware	10.00	0.64	09(02,32)	10.20	North Carolina	8.75	0.19	21(14,33)	-3.57
D.C.	14.97	1.08	01(01,02)	65.04 ^c	North Dakota	5.40	0.55	51(42,51)	-40.42 ^c
Florida	8.73	0.12	22(16,30)	-3.73	Ohio	8.20	0.17	32(23,38)	-9.65
Georgia	8.63	0.21	23(15,35)	-4.90	Oklahoma	8.87	0.30	20(10,35)	-2.26
Hawaii	11.31	0.55	04(02,12)	24.73 ^c	Oregon	9.29	0.30	16(08,29)	2.45
Idaho	6.46	0.41	47(36,51)	-28.73 ^c	Pennsylvania	8.49	0.15	27(19,35)	-6.36
Illinois	8.52	0.17	24(18,35)	-6.03	Rhode Island	10.57	0.62	06(02,21)	16.54 ^c
Indiana	7.69	0.22	38(28,45)	-15.22 ^c	South Carolina	8.98	0.28	19(10,33)	-0.97
Iowa	6.88	0.29	46(36,50)	-24.13 ^c	South Dakota	5.96	0.52	50(38,51)	-34.29 ^c
Kansas	7.18	0.31	43(32,48)	-20.80 ^c	Tennessee	9.71	0.25	12(06,20)	7.02
Kentucky	8.41	0.28	31(15,39)	-7.33	Texas	11.54	0.15	03(02,05)	27.24 ^c
Louisiana	11.81	0.33	02(01,06)	30.21 ^c	Utah	5.97	0.34	49(44,51)	-34.18 ^c
Maine	7.49	0.44	39(21,48)	-17.37 ^c	Vermont	7.81	0.68	35(10,50)	-13.89
Maryland	9.40	0.26	13(07,25)	3.59	Virginia	8.43	0.21	29(17,37)	-7.03
Massachusetts	10.07	0.25	08(05,16)	11.05	Washington	9.36	0.24	15(08,24)	3.23
Michigan	8.07	0.18	33(24,40)	-11.06	West Virginia	7.29	0.37	41(27,49)	-19.63 ^c
Minnesota	7.24	0.23	42(34,48)	-20.16 ^c	Wisconsin	7.48	0.22	40(32,47)	-17.54 ^c
Mississippi	10.43	0.39	07(03,16)	15.01 ^c	Wyoming	6.96	0.70	45(16,51)	-23.24 ^c
Missouri	9.09	0.24	17(10,29)	0.19					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.
 - Statistic not shown. Rate based on less than 16 cases for the time interval.