


Table 14.20
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.65	0.02		
High Five States				
D.C.	5.35	0.56	01(01,16)	46.70 ^c
Texas	4.58	0.09	02(01,07)	25.42 ^c
Louisiana	4.57	0.19	03(01,09)	25.10 ^c
California	4.49	0.07	04(01,07)	22.96 ^c
Hawaii	4.39	0.31	05(01,23)	20.29 ^c
Low Five States				
Nebraska	2.91	0.23	47(16,51)	-20.19 ^c
Idaho	2.91	0.25	48(16,51)	-20.37 ^c
Vermont	2.88	0.38	49(07,51)	-21.08 ^c
New Hampshire	2.58	0.25	50(29,51)	-29.41 ^c
North Dakota	2.23	0.31	51(35,51)	-38.83 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.85	0.16	10(05,30)	5.43	Montana	3.08	0.31	41(08,51)	-15.60 ^c
Alaska	3.33	0.51	32(01,51)	-8.86	Nebraska	2.91	0.23	47(16,51)	-20.19 ^c
Arizona	3.85	0.14	09(06,26)	5.60	Nevada	3.54	0.22	21(06,47)	-3.06
Arkansas	3.82	0.20	11(04,36)	4.53	New Hampshire	2.58	0.25	50(29,51)	-29.41 ^c
California	4.49	0.07	04(01,07)	22.96 ^c	New Jersey	3.38	0.11	29(14,43)	-7.47
Colorado	3.06	0.15	42(22,50)	-16.18 ^c	New Mexico	4.06	0.26	08(01,33)	11.37
Connecticut	2.96	0.16	44(23,51)	-18.96 ^c	New York	3.59	0.08	18(10,31)	-1.60
Delaware	3.77	0.36	12(01,49)	3.38	North Carolina	3.21	0.11	39(21,47)	-12.04
D.C.	5.35	0.56	01(01,16)	46.70 ^c	North Dakota	2.23	0.31	51(35,51)	-38.83 ^c
Florida	3.34	0.07	31(19,42)	-8.42	Ohio	3.24	0.09	37(21,45)	-11.24
Georgia	3.41	0.12	27(12,42)	-6.67	Oklahoma	3.55	0.18	20(08,43)	-2.64
Hawaii	4.39	0.31	05(01,23)	20.29 ^c	Oregon	3.68	0.17	16(06,39)	0.84
Idaho	2.91	0.25	48(16,51)	-20.37 ^c	Pennsylvania	3.35	0.09	30(17,42)	-8.31
Illinois	3.69	0.10	14(08,28)	1.23	Rhode Island	3.62	0.33	17(02,49)	-0.69
Indiana	3.46	0.13	25(10,43)	-5.28	South Carolina	3.28	0.15	34(14,48)	-10.17
Iowa	2.93	0.17	46(25,51)	-19.85 ^c	South Dakota	3.32	0.36	33(04,51)	-9.08
Kansas	3.27	0.20	35(11,49)	-10.36	Tennessee	3.69	0.14	15(07,35)	1.05
Kentucky	3.59	0.16	19(07,41)	-1.63	Texas	4.58	0.09	02(01,07)	25.42 ^c
Louisiana	4.57	0.19	03(01,09)	25.10 ^c	Utah	3.22	0.24	38(08,50)	-11.67
Maine	3.19	0.26	40(08,50)	-12.63	Vermont	2.88	0.38	49(07,51)	-21.08 ^c
Maryland	3.71	0.15	13(07,35)	1.54	Virginia	3.24	0.12	36(18,47)	-11.13
Massachusetts	3.53	0.13	22(09,41)	-3.16	Washington	4.07	0.15	07(03,19)	11.42
Michigan	3.42	0.10	26(13,42)	-6.30	West Virginia	2.95	0.22	45(17,51)	-19.27 ^c
Minnesota	3.05	0.14	43(22,50)	-16.31 ^c	Wisconsin	3.40	0.14	28(12,45)	-6.95
Mississippi	4.37	0.22	06(01,16)	19.72 ^c	Wyoming	3.48	0.48	24(01,51)	-4.61
Missouri	3.50	0.14	23(10,42)	-4.10					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.