

Table 14.21
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	38,940	9,558	3,239	1,214	716	527	53,480	54,464	490	54,954
	Males	28,968	6,762	2,047	722	410	213	38,824	39,263	71	39,334
	Females	9,972	2,796	1,192	492	306	314	14,656	15,201	419	15,620
White ^b	Both Sexes	28,607	6,931	2,307	860	559	438	39,117	39,907	401	40,308
	Males	21,328	5,056	1,433	495	304	159	28,559	28,887	264 ⁱ	29,151
	Females	7,279	1,875	874	365	255	279	10,558	11,020	137 ⁱ	11,157
Black ^b	Both Sexes	5,786	983	231	94	71	23	7,143	7,237	+	+
	Males	4,360	559	132	47	35	13	5,122	5,171	+	+
	Females	1,426	424	99	47	36	10	2,021	2,066	+	+
Asian/ Pacific Islander ^c	Both Sexes	3,850	1,443	626	+	+	+	6,121	+	+	+
	Males	2,716	1,039	422	+	+	+	4,323	+	+	+
	Females	1,134	404	204	+	+	+	1,798	+	+	+
Hispanic ^d	Both Sexes	6,209	1,333	497	+	+	+	8,291	+	+	+
	Males	4,545	948	284	+	+	+	5,953	+	+	+
	Females	1,664	385	213	+	+	+	2,338	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0191%	0.0029%	0.0027%	0.0013%	0.0019%	0.0069%	0.0358%	0.0731%	0.0703%	0.0447%	0.0171%
	Males	0.0278%	0.0033%	0.0026%	0.0015%	0.0023%	0.0096%	0.0571%	0.1173%	0.1004%	0.0729%	0.0259%
	Females	0.0107%	0.0024%	0.0028%	0.0011%	0.0015%	0.0042%	0.0154%	0.0331%	0.0457%	0.0279%	0.0095%
White ^c	Both Sexes	0.0161%	0.0032%	0.0028%	0.0015%	0.0014%	0.0054%	0.0309%	0.0609%	0.0488%	0.0283%	0.0137%
	Males	0.0234%	0.0036%	0.0027%	0.0016%	0.0013%	0.0072%	0.0491%	0.0977%	0.0702%	0.0462%	0.0204%
	Females	0.0087%	0.0029%	0.0030%	0.0013%	0.0014%	0.0035%	0.0128%	0.0265%	0.0309%	0.0175%	0.0075%
Black ^c	Both Sexes	0.0158%	0.0009%	0.0007%	0.0014%	0.0024%	0.0046%	0.0338%	0.0959%	0.0554%	0.0313%	0.0163%
	Males	0.0232%	-	-	0.0017%	0.0036%	0.0057%	0.0513%	0.1628%	0.0820%	0.0560%	0.0258%
	Females	0.0091%	-	-	-	-	0.0036%	0.0185%	0.0437%	0.0368%	0.0191%	0.0090%
Asian/ Pacific Islander ^c	Both Sexes	0.0377%	0.0027%	0.0037%	0.0007%	0.0039%	0.0151%	0.0630%	0.1235%	0.1978%	0.1499%	0.0366%
	Males	0.0552%	0.0037%	0.0036%	-	0.0059%	0.0240%	0.1079%	0.2002%	0.2856%	0.2405%	0.0570%
	Females	0.0217%	0.0016%	0.0039%	-	0.0021%	0.0072%	0.0244%	0.0605%	0.1296%	0.0945%	0.0203%
Hispanic ^d	Both Sexes	0.0159%	0.0038%	0.0026%	0.0012%	0.0010%	0.0073%	0.0508%	0.0980%	0.0935%	0.0683%	0.0229%
	Males	0.0225%	0.0046%	0.0027%	0.0012%	0.0009%	0.0116%	0.0852%	0.1560%	0.1269%	0.1039%	0.0349%
	Females	0.0091%	0.0030%	0.0025%	0.0011%	0.0011%	0.0027%	0.0177%	0.0481%	0.0688%	0.0468%	0.0127%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <38 and >=38. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.