Table 15.21 Lung and Bronchus (Invasive)

SEER Incidence and U.S. Mortality Age-Adjusted Rates and Trends^a By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas ^b Rate 2009-2013 Rate per 100,000 persons			SEER 13 Areas ^b Trend 2004-2013 ^c AAPC (%)			SEER 9 Areas ^b Trend 2004-2013 ^d AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
RACE/ETHNICITY									
All Races	57.3	67.9	49.4	-2.1*	-2.9*	-1.5*	-1.8*	-2.6*	-1.2*
White White Hispanic ^e White Non-Hispanic ^e	58.8 30.7 63.2	68.1 37.9 72.7	51.8 25.6 56.2	-2.1* -1.3* -2.0*		-0.4*	-1.8* -	-2.6* -	-1.2* -
WHITE NOH-HISPAHIC	63.2	12.1	50.2	-2.0"	-2.7"	-1.4"	_	_	_
Black	65.0	87.3	50.0	-2.7*	-2.7*	-1.7*	-1.9*	-2.8*	-1.3*
Asian/Pacific Islander	36.7	47.9	28.5	-1.4*	-2.2*	0.2	-	-	-
Amer Ind/Alaska Natf	38.9	48.6	32.0	-0.9*	-1.6*	-3.5	-	-	-
Hispanic ^e	29.7	36.6	24.8	-1.3*	-2.1*	-0.4*	-	-	-
U.S. Mortality ^g	Rate 2009-2013 Rate per 100,000 persons			Trend 2004-2013 ^h AAPC (%)					
	Total	Males	Females	Total	Males	Females			
RACE/ETHNICITY									
All Races	46.0	57.8	37.0	-2.3*	-2.9*	-1.6*			
White	46.7	57.7	38.3	-2.1*					
White Hispanic ^e White Non-Hispanic ^e	20.7	29.7 60.2	14.2 40.6	-2.3* -2.0*	-3.0* -2.7*				
white Non-Hispanic	49.1	60.2	40.6	-2.0^	-2./*	-1.4			
Black	49.4	70.6	35.3	-2.7*	-3.4*	-1.9*			
Asian/Pacific Islander	24.2	32.7	18.1	-1.2*	-2.2*	-0.4*			
Amer Ind/Alaska Nat Total U.S. CHSDA Counties	30.3 37.8	37.5 46.6	24.9 31.3	-1.2* -0.3	-2.1* -1.1*				
Non-CHSDA Counties	21.7	27.1	17.5	-2.3*	-3.3*	-2.2*			
Hispanic ^e	19.7	28.3	13.5	-2.5*	-3.3*	-1.4*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, b Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and $\bar{\text{R}}\text{ural Georgia.}$

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky,

- Louisiana, New Jersey, and Georgia excluding ATT/RG.
 The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.
- The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over d diagnosis years 1975-2013.
- Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.
- Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- US Mortality Files, National Center for Health Statistics, CDC.
- The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013. The APC is significantly different from zero (p<.05).