


Table 15.24
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	45.97	0.05		
High Five States				
Kentucky	69.96	0.54	01(01,01)	52.19 ^c
Arkansas	61.86	0.60	02(02,04)	34.57 ^c
West Virginia	61.61	0.72	03(02,05)	34.03 ^c
Mississippi	60.00	0.62	04(02,06)	30.51 ^c
Tennessee	59.95	0.41	05(03,05)	30.41 ^c
Low Five States				
California	34.81	0.14	47(47,47)	-24.27 ^c
Colorado	33.11	0.38	48(48,50)	-27.98 ^c
Hawaii	32.60	0.64	49(48,50)	-29.08 ^c
New Mexico	32.12	0.54	50(48,50)	-30.13 ^c
Utah	20.35	0.44	51(51,51)	-55.74 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	57.29	0.46	07(06,10)	24.63 ^c	Montana	40.83	0.83	39(35,44)	-11.18 ^c
Alaska	50.78	1.52	17(11,27)	10.46 ^c	Nebraska	43.70	0.66	34(29,36)	-4.94
Arizona	37.46	0.32	46(44,46)	-18.50 ^c	Nevada	47.77	0.60	21(18,27)	3.92
Arkansas	61.86	0.60	02(02,04)	34.57 ^c	New Hampshire	47.54	0.80	23(17,29)	3.42
California	34.81	0.14	47(47,47)	-24.27 ^c	New Jersey	40.86	0.29	38(37,43)	-11.10 ^c
Colorado	33.11	0.38	48(48,50)	-27.98 ^c	New Mexico	32.12	0.54	50(48,50)	-30.13 ^c
Connecticut	40.63	0.44	41(36,44)	-11.62 ^c	New York	41.20	0.20	37(36,42)	-10.37 ^c
Delaware	52.19	0.99	13(11,18)	13.53 ^c	North Carolina	51.79	0.32	15(12,17)	12.66 ^c
D.C.	40.64	1.20	40(34,45)	-11.58 ^c	North Dakota	40.25	1.01	43(35,45)	-12.44 ^c
Florida	45.25	0.19	30(27,32)	-1.56	Ohio	54.26	0.29	11(11,13)	18.04 ^c
Georgia	48.96	0.33	19(17,22)	6.51 ^c	Oklahoma	57.87	0.53	06(05,08)	25.89 ^c
Hawaii	32.60	0.64	49(48,50)	-29.08 ^c	Oregon	45.72	0.46	28(25,33)	-0.53
Idaho	38.00	0.69	45(42,46)	-17.34 ^c	Pennsylvania	47.49	0.25	24(20,27)	3.31
Illinois	48.47	0.27	20(17,23)	5.43 ^c	Rhode Island	49.00	0.90	18(15,27)	6.60 ^c
Indiana	56.28	0.40	09(07,10)	22.42 ^c	South Carolina	52.01	0.45	14(12,17)	13.15 ^c
Iowa	47.15	0.51	26(20,28)	2.57	South Dakota	43.58	0.96	35(28,39)	-5.19 ^c
Kansas	47.41	0.55	25(18,28)	3.13	Tennessee	59.95	0.41	05(03,05)	30.41 ^c
Kentucky	69.96	0.54	01(01,01)	52.19 ^c	Texas	41.97	0.20	36(35,38)	-8.69 ^c
Louisiana	56.37	0.49	08(06,10)	22.62 ^c	Utah	20.35	0.44	51(51,51)	-55.74 ^c
Maine	53.71	0.79	12(10,15)	16.85 ^c	Vermont	47.55	1.13	22(17,32)	3.45
Maryland	44.55	0.39	32(28,35)	-3.09	Virginia	46.97	0.34	27(21,28)	2.18
Massachusetts	45.31	0.35	29(27,33)	-1.44	Washington	44.01	0.36	33(30,35)	-4.26
Michigan	50.93	0.30	16(14,18)	10.80 ^c	West Virginia	61.61	0.72	03(02,05)	34.03 ^c
Minnesota	40.40	0.38	42(37,44)	-12.11 ^c	Wisconsin	44.92	0.37	31(28,34)	-2.29
Mississippi	60.00	0.62	04(02,06)	30.51 ^c	Wyoming	38.53	1.15	44(37,46)	-16.19 ^c
Missouri	55.99	0.40	10(07,11)	21.80 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 5% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.