

Table 18.9
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Survival Time																						
1-year	68.0	69.5	72.3	71.9	73.9	70.6	74.3	74.9	73.6	72.4	72.6	73.5	77.2	74.6	76.0	74.0	76.9	77.6	81.8	81.3	79.4	80.9
2-year	52.1	54.5	57.6	57.4	58.8	57.6	60.6	62.3	59.8	59.9	59.6	62.7	64.9	66.2	64.6	66.1	66.4	68.6	72.4	71.8	71.0	
3-year	39.6	42.8	45.3	46.2	47.9	44.9	48.7	51.1	47.9	50.0	49.4	53.3	55.7	58.4	57.2	57.9	57.6	61.3	64.8	65.0		
4-year	31.5	34.2	35.7	36.9	40.1	38.2	38.7	39.8	38.8	41.3	43.9	46.6	47.6	50.4	50.4	50.9	49.9	55.3	58.9			
5-year	25.1	26.9	27.5	29.9	33.4	31.8	31.3	34.2	31.3	34.5	37.3	39.7	41.9	44.9	46.1	45.8	45.6	49.6				
6-year	20.1	21.0	22.1	24.5	28.3	27.1	26.3	29.7	27.2	29.2	32.5	35.0	37.5	39.5	41.1	40.6	40.5					
7-year	16.2	16.7	17.5	20.4	23.7	23.7	22.5	24.4	23.5	25.4	29.0	31.4	32.9	35.1	36.2	35.8						
8-year	13.5	13.4	14.9	17.3	20.4	21.4	19.8	22.2	20.2	22.7	25.5	28.2	29.7	31.3	33.1							
9-year	11.5	11.2	11.9	15.3	18.3	19.5	17.1	19.5	18.4	20.5	22.9	25.3	26.7	29.4								
10-year	9.8	9.6	10.4	13.7	17.2	17.2	15.5	17.8	16.4	19.4	20.9	22.5	23.7									
11-year	8.7	8.6	9.2	12.6	15.6	15.6	14.3	16.7	14.2	18.4	19.5	19.5										
12-year	7.8	7.6	8.4	11.1	14.3	14.8	12.9	14.9	13.7	17.0	18.5											
13-year	6.7	6.6	7.6	10.2	13.5	13.3	11.7	14.0	12.7	15.9												
14-year	6.1	6.0	7.0	9.5	12.4	12.6	11.0	13.7	11.4													
15-year	5.3	5.7	6.1	9.2	11.9	11.4	10.8	12.1														
16-year	4.9	5.1	5.7	8.6	10.8	9.7	10.3															
17-year	4.6	4.7	5.4	8.1	10.0	8.8																
18-year	4.0	4.3	5.0	7.5	9.8																	
19-year	3.8	3.8	4.9	6.8																		
20-year	3.3	3.5	4.7																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).