

Table 18.10  
MyelomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Survival Time																						
1-year	67.8	69.6	73.2	72.5	74.2	71.2	74.1	76.2	75.6	73.6	74.2	74.2	76.9	76.6	76.1	73.7	77.5	76.6	83.6	83.0	79.8	79.8
2-year	52.3	54.7	59.3	58.4	59.8	58.3	61.0	63.6	61.3	61.6	61.5	61.4	65.3	67.9	64.0	66.7	68.0	67.5	73.6	73.5	71.8	
3-year	39.0	42.8	46.2	46.6	49.2	45.0	50.3	53.4	50.7	52.5	50.8	52.4	56.3	60.1	56.4	58.1	59.9	58.8	66.1	66.0		
4-year	30.8	33.6	36.6	37.0	41.2	39.8	40.8	43.6	42.7	43.2	45.8	46.6	47.7	51.3	50.2	51.6	51.7	52.2	60.5			
5-year	24.3	25.8	28.4	30.3	36.2	33.7	31.5	38.0	34.6	35.8	39.7	41.4	42.2	45.8	46.7	47.3	47.5	47.0				
6-year	19.2	20.8	23.5	25.2	31.6	29.7	26.4	33.9	29.8	30.0	34.8	36.1	37.7	39.9	41.0	42.8	42.4					
7-year	15.4	16.9	19.1	21.3	26.0	25.8	22.3	28.0	25.9	26.0	31.0	32.7	34.2	35.6	35.5	37.3						
8-year	13.0	13.8	16.2	18.6	22.0	23.9	20.1	25.5	21.5	23.3	27.0	29.4	30.6	32.2	33.4							
9-year	11.0	11.7	12.9	16.7	20.0	21.2	18.1	23.3	19.3	19.7	24.3	26.3	28.2	30.3								
10-year	9.2	9.9	11.2	14.8	19.0	18.3	16.3	21.6	17.3	18.4	21.6	23.6	23.5									
11-year	7.5	8.9	9.9	13.6	17.3	17.0	14.8	20.7	14.2	17.0	20.4	20.9										
12-year	6.9	7.7	9.4	12.0	15.4	16.6	14.3	18.6	13.5	15.5	19.2											
13-year	5.8	6.7	8.6	11.2	14.8	14.4	13.0	17.1	12.7	14.6												
14-year	5.2	6.1	8.0	10.4	13.1	14.0	12.6	16.7	11.1													
15-year	4.6	5.7	7.0	10.0	12.9	12.6	12.1	15.8														
16-year	4.4	5.2	6.7	9.6	11.1	9.5	11.6															
17-year	3.9	4.8	6.3	8.9	10.3	9.2																
18-year	3.5	4.4	6.1	8.1	9.9																	
19-year	3.3	4.2	6.1	7.2																		
20-year	2.6	3.9	5.9																			

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).