

Table 18.11
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Survival Time																						
1-year	68.2	69.3	71.2	71.1	73.6	69.9	74.6	73.4	71.3	71.0	70.8	72.6	77.5	72.2	75.9	74.4	76.2	78.7	79.7	79.0	78.9	82.4
2-year	52.0	54.3	55.9	56.3	57.8	56.8	60.1	60.9	58.1	57.9	57.6	64.4	64.4	64.2	65.3	65.5	64.2	69.8	70.9	69.7	69.9	
3-year	40.4	42.9	44.3	45.7	46.3	44.9	46.9	48.5	44.8	47.1	47.7	54.5	54.9	56.3	58.0	57.7	54.3	63.9	63.2	63.8		
4-year	32.2	34.8	34.7	36.8	39.0	36.4	36.5	35.6	34.4	39.0	41.7	46.7	47.5	49.4	50.5	50.0	47.5	58.6	57.1			
5-year	26.1	28.0	26.5	29.4	30.2	29.8	31.1	29.9	27.7	32.9	34.6	37.5	41.5	44.0	45.4	44.3	42.9	52.4				
6-year	21.0	21.2	20.7	23.7	24.7	24.4	26.3	25.2	24.2	28.3	29.9	33.6	37.2	38.9	41.2	38.2	37.8					
7-year	17.1	16.4	15.9	19.4	21.3	21.5	22.6	20.4	20.7	24.5	26.7	29.7	31.4	34.5	37.0	34.3						
8-year	14.1	13.0	13.7	15.9	18.5	18.6	19.4	18.5	18.6	21.9	23.7	26.7	28.6	30.1	32.6							
9-year	12.0	10.7	10.9	13.9	16.3	17.7	16.1	15.4	17.2	21.5	21.3	24.0	25.1	28.3								
10-year	10.5	9.4	9.6	12.4	15.1	15.9	14.5	13.8	15.4	20.5	20.0	21.2	23.9									
11-year	10.0	8.4	8.4	11.6	13.7	14.1	13.6	12.3	14.1	20.0	18.5	17.6										
12-year	8.7	7.5	7.3	10.1	12.8	13.0	11.4	10.9	13.9	18.5	17.7											
13-year	7.5	6.5	6.7	9.2	12.0	11.9	10.2	10.6	12.6	17.5												
14-year	6.9	6.0	6.0	8.5	11.4	11.0	9.4	10.4	11.7													
15-year	6.0	5.6	5.2	8.2	10.8	10.2	9.4	8.0														
16-year	5.4	5.0	4.7	7.5	10.3	9.8	8.9															
17-year	5.2	4.5	4.5	7.2	9.5	8.3																
18-year	4.5	4.1	4.0	6.7	9.5																	
19-year	4.3	3.4	3.8	6.4																		
20-year	3.9	3.1	3.5																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).