

Table 18.13

Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.88	0.48	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.64	0.33
	10	0.00	0.00	0.01	0.89	0.48		10	0.00	0.00	0.01	0.65	0.33
	20	0.00	0.01	0.04	0.89	0.48		20	0.00	0.01	0.03	0.65	0.33
	30	0.01	0.04	0.14	0.90	0.49		30	0.01	0.03	0.08	0.65	0.34
	40	0.04	0.13	0.34	0.91	0.50		40	0.02	0.08	0.20	0.65	0.34
	50	0.10	0.31	0.63	0.90	0.51		50	0.06	0.18	0.39	0.64	0.34
	60	0.23	0.57	0.81	0.86	0.51		60	0.13	0.34	0.53	0.61	0.33
	70	0.40	0.68	-	0.74	0.49		70	0.23	0.44	-	0.52	0.31
White	80	0.40	-	-	0.48	0.39	American Indian/ Alaska Native ^a	80	0.25	-	-	0.36	0.23
	0	0.00	0.00	0.00	0.82	0.46		0	0.00	0.00	0.00	0.73	0.33
	10	0.00	0.00	0.01	0.83	0.46		10	0.00	0.00	0.01	0.74	0.34
	20	0.00	0.01	0.04	0.84	0.47		20	0.00	0.00	0.04	0.74	0.34
	30	0.01	0.04	0.12	0.85	0.47		30	0.00	0.04	0.05	0.76	0.35
	40	0.03	0.12	0.30	0.85	0.48		40	0.04	0.05	0.25	0.78	0.36
	50	0.09	0.28	0.59	0.85	0.49		50	0.02	0.22	0.45	0.78	0.38
	60	0.21	0.54	0.77	0.82	0.49		60	0.22	0.47	0.69	0.83	0.39
Black	70	0.38	0.65	-	0.71	0.47	Hispanic ^b	70	0.30	0.56	-	0.72	0.39
	80	0.39	-	-	0.46	0.38		80	0.38	-	-	0.61	0.36
	0	0.00	0.00	0.00	1.38	0.68		0	0.00	0.00	0.00	0.91	0.46
	10	0.00	0.00	0.02	1.41	0.70		10	0.00	0.00	0.01	0.91	0.46
	20	0.00	0.02	0.09	1.42	0.70		20	0.00	0.01	0.04	0.92	0.46
	30	0.02	0.09	0.28	1.45	0.72		30	0.01	0.04	0.13	0.93	0.47
	40	0.07	0.26	0.67	1.46	0.73		40	0.03	0.12	0.32	0.93	0.47
	50	0.20	0.62	1.12	1.45	0.75		50	0.10	0.29	0.62	0.92	0.48
	60	0.48	1.03	1.35	1.41	0.76		60	0.21	0.55	0.82	0.87	0.47
	70	0.71	1.11	-	1.18	0.73		70	0.39	0.69	-	0.75	0.44
	80	0.65	-	-	0.76	0.61		80	0.41	-	-	0.48	0.32

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.