

Table 18.14

Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Females, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.65	0.39	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.46	0.25
	10	0.00	0.00	0.01	0.65	0.39		10	0.00	0.00	0.00	0.46	0.25
	20	0.00	0.01	0.03	0.65	0.39		20	0.00	0.00	0.02	0.46	0.25
	30	0.01	0.03	0.10	0.65	0.39		30	0.00	0.02	0.05	0.46	0.25
	40	0.03	0.10	0.25	0.65	0.40		40	0.01	0.05	0.14	0.46	0.25
	50	0.07	0.23	0.44	0.64	0.40		50	0.04	0.13	0.28	0.45	0.25
	60	0.16	0.39	0.56	0.59	0.40		60	0.10	0.25	0.37	0.43	0.25
	70	0.25	0.44	-	0.48	0.36		70	0.16	0.28	-	0.34	0.23
White	80	0.24	-	-	0.29	0.28	American Indian/ Alaska Native ^a	80	0.14	-	-	0.21	0.17
	0	0.00	0.00	0.00	0.57	0.35		0	0.00	0.00	0.00	0.79	0.30
	10	0.00	0.00	0.01	0.57	0.36		10	0.00	0.00	0.00	0.80	0.30
	20	0.00	0.01	0.02	0.57	0.36		20	0.00	0.00	0.03	0.80	0.30
	30	0.00	0.02	0.08	0.57	0.36		30	0.00	0.03	0.09	0.81	0.31
	40	0.02	0.08	0.21	0.57	0.36		40	0.03	0.09	0.28	0.82	0.31
	50	0.06	0.19	0.38	0.56	0.37		50	0.07	0.27	0.47	0.82	0.32
	60	0.14	0.34	0.49	0.53	0.36		60	0.21	0.42	0.74	0.79	0.32
Black	70	0.22	0.39	-	0.43	0.34	Hispanic ^b	70	0.23	0.59	-	0.65	0.27
	80	0.22	-	-	0.27	0.26		80	0.46	-	-	0.54	0.16
	0	0.00	0.00	0.00	1.28	0.69		0	0.00	0.00	0.00	0.68	0.39
	10	0.00	0.00	0.01	1.30	0.71		10	0.00	0.00	0.01	0.68	0.39
	20	0.00	0.01	0.08	1.30	0.71		20	0.00	0.01	0.03	0.69	0.40
	30	0.01	0.08	0.25	1.31	0.71		30	0.00	0.03	0.09	0.69	0.40
	40	0.06	0.24	0.57	1.32	0.72		40	0.02	0.09	0.24	0.69	0.40
	50	0.18	0.52	0.95	1.29	0.73		50	0.06	0.22	0.44	0.67	0.40
	60	0.36	0.83	1.13	1.20	0.73		60	0.16	0.39	0.57	0.62	0.39
	70	0.53	0.89	-	0.96	0.67		70	0.25	0.44	-	0.50	0.35
	80	0.49	-	-	0.58	0.53		80	0.23	-	-	0.30	0.26

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.