

Table 18.15  
Myeloma

SEER Incidence and U.S. Mortality  
Age-Adjusted Rates and Trends<sup>a</sup>  
By Race/Ethnicity and Sex

SEER Incidence	SEER 18 Areas <sup>b</sup> Rate 2009-2013 Rate per 100,000 persons			SEER 13 Areas <sup>b</sup> Trend 2004-2013 <sup>c</sup> AAPC (%)			SEER 9 Areas <sup>b</sup> Trend 2004-2013 <sup>d</sup> AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
	RACE/ETHNICITY								
All Races	6.5	8.2	5.2	0.8	0.8*	0.4*	0.8*	0.9*	0.7*
White	5.9	7.7	4.5	0.7*	0.8*	0.3	0.8*	0.9*	0.5*
White Hispanic <sup>e</sup>	6.1	7.7	4.9	0.2	0.4	-0.1	-	-	-
White Non-Hispanic <sup>e</sup>	5.9	7.7	4.4	0.7*	0.8*	0.3	-	-	-
Black	13.2	15.7	11.5	0.8*	1.1*	0.4	0.9*	0.9*	0.9*
Asian/Pacific Islander	3.7	4.7	3.0	0.6	0.7	0.6	-	-	-
Amer Ind/Alaska Nat <sup>f</sup>	5.0	5.2	4.8	0.2	-0.2	0.2	-	-	-
Hispanic <sup>e</sup>	6.0	7.5	4.9	0.2	0.4	-0.1	-	-	-
<u>U.S. Mortality<sup>g</sup></u>									
	Rate 2009-2013 Rate per 100,000 persons			Trend 2004-2013 <sup>h</sup> AAPC (%)					
	Total	Males	Females	Total	Males	Females			
	RACE/ETHNICITY								
All Races	3.3	4.2	2.7	-0.9*	-1.0*	-1.0*			
White	3.1	4.0	2.4	-0.9*	-0.8*	-1.2*			
White Hispanic <sup>e</sup>	2.9	3.6	2.4	-0.6*	-0.6	-0.7*			
White Non-Hispanic <sup>e</sup>	3.1	4.0	2.4	-0.8*	-0.9*	-1.2*			
Black	6.2	7.5	5.4	-0.8	-1.4*	-0.5			
Asian/Pacific Islander	1.6	2.2	1.3	-0.8*	-0.3	-1.2*			
Amer Ind/Alaska Nat									
Total U.S.	2.2	2.5	2.0	-2.1*	-2.0*	-2.5*			
CHSDA Counties	2.9	3.4	2.6	-1.8*	-1.3	-2.3			
Non-CHSDA Counties	1.4	1.6	1.2	-1.7	-2.7	-			
Hispanic <sup>e</sup>	2.8	3.5	2.3	-0.7*	-0.7*	-0.8*			

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 4.3.0.0, April 2016, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

<sup>a</sup> Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry, and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG.

<sup>c</sup> The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1992-2013.

<sup>d</sup> The 2004-2013 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2013.

<sup>e</sup> Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The 2004-2013 Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

<sup>f</sup> Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

<sup>g</sup> US Mortality Files, National Center for Health Statistics, CDC.

<sup>h</sup> The 2004-2013 mortality AAPCs are based on a Joinpoint analysis using years of death 1992-2013.

\* The APC is significantly different from zero (p<.05).