


Table 18.20
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013


Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.69	0.02		
High Five States				
D.C.	3.52	0.45	01(01,47)	30.56 ^c
Maryland	3.37	0.14	02(01,14)	25.07 ^c
Mississippi	3.24	0.19	03(01,30)	20.28 ^c
South Carolina	3.19	0.15	04(01,24)	18.47 ^c
Alabama	3.18	0.14	05(01,23)	18.02 ^c
Low Five States				
Nevada	2.29	0.18	47(17,51)	-15.16 ^c
Rhode Island	2.27	0.25	48(10,51)	-15.80 ^c
Vermont	2.24	0.32	49(03,51)	-16.80 ^c
New Mexico	2.17	0.19	50(22,51)	-19.41 ^c
Hawaii	1.78	0.20	51(41,51)	-34.04 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.18	0.14	05(01,23)	18.02 ^c	Montana	2.81	0.29	19(01,50)	4.16
Alaska	2.69	0.48	30(01,51)	-0.22	Nebraska	2.56	0.21	36(05,50)	-4.89
Arizona	2.30	0.11	46(30,51)	-14.63	Nevada	2.29	0.18	47(17,51)	-15.16 ^c
Arkansas	2.72	0.17	24(04,48)	1.07	New Hampshire	2.52	0.25	42(04,51)	-6.58
California	2.46	0.05	43(30,48)	-8.74	New Jersey	2.52	0.09	39(19,48)	-6.31
Colorado	2.41	0.14	45(19,50)	-10.50	New Mexico	2.17	0.19	50(22,51)	-19.41 ^c
Connecticut	2.52	0.14	41(13,50)	-6.49	New York	2.56	0.06	37(22,46)	-4.97
Delaware	2.84	0.31	17(01,50)	5.22	North Carolina	3.13	0.10	06(01,21)	16.16 ^c
D.C.	3.52	0.45	01(01,47)	30.56 ^c	North Dakota	2.78	0.36	20(01,51)	3.08
Florida	2.42	0.06	44(30,49)	-10.37	Ohio	2.98	0.09	11(03,26)	10.50
Georgia	3.08	0.11	07(01,25)	14.18	Oklahoma	2.58	0.15	35(10,50)	-4.37
Hawaii	1.78	0.20	51(41,51)	-34.04 ^c	Oregon	2.62	0.15	31(08,49)	-2.86
Idaho	2.61	0.25	32(02,51)	-3.00	Pennsylvania	2.72	0.08	26(12,40)	0.93
Illinois	2.72	0.08	25(11,41)	0.99	Rhode Island	2.27	0.25	48(10,51)	-15.80 ^c
Indiana	2.83	0.12	18(05,40)	4.87	South Carolina	3.19	0.15	04(01,24)	18.47 ^c
Iowa	2.59	0.16	33(08,49)	-3.95	South Dakota	3.06	0.33	08(01,49)	13.54
Kansas	2.90	0.18	15(01,45)	7.44	Tennessee	2.94	0.12	13(03,33)	9.20
Kentucky	2.72	0.14	27(05,47)	0.80	Texas	2.54	0.06	38(23,46)	-5.72
Louisiana	3.04	0.15	09(01,35)	13.01	Utah	2.78	0.22	21(01,49)	2.99
Maine	2.59	0.23	34(04,50)	-4.05	Vermont	2.24	0.32	49(03,51)	-16.80 ^c
Maryland	3.37	0.14	02(01,14)	25.07 ^c	Virginia	2.99	0.11	10(02,31)	10.88
Massachusetts	2.70	0.11	29(10,45)	0.39	Washington	2.71	0.12	28(08,45)	0.72
Michigan	2.87	0.10	16(06,34)	6.66	West Virginia	2.92	0.21	14(01,46)	8.49
Minnesota	2.52	0.13	40(15,49)	-6.39	Wisconsin	2.74	0.12	23(06,44)	1.82
Mississippi	3.24	0.19	03(01,30)	20.28 ^c	Wyoming	2.96	0.44	12(01,51)	9.89
Missouri	2.76	0.12	22(07,44)	2.40					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.