

Table 19.10
Non-Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Survival Time																						
1-year	69.4	72.7	68.9	67.1	66.6	71.2	74.7	74.9	75.8	76.5	77.6	80.6	79.6	80.1	81.0	81.0	81.4	83.5	82.7	83.6	83.6	83.8
2-year	59.4	63.7	60.5	58.2	58.3	63.6	67.1	68.2	69.1	70.3	71.2	75.2	74.8	75.4	76.1	76.5	77.8	77.7	78.0	78.6	79.5	
3-year	53.4	58.0	55.7	53.7	54.5	59.8	63.8	64.5	64.4	66.7	67.9	72.0	72.5	72.6	73.0	73.6	75.3	75.0	75.5	76.3		
4-year	49.3	53.9	51.6	50.4	51.5	56.1	61.0	61.9	61.5	63.9	65.6	70.0	70.8	70.0	70.9	71.4	73.3	73.1	73.4			
5-year	45.4	50.1	48.2	47.6	48.5	53.6	58.2	59.1	59.0	61.7	63.5	67.9	68.8	68.7	69.7	68.9	70.9	71.9				
6-year	42.6	46.6	45.5	45.5	46.8	51.0	56.1	56.4	56.2	60.2	61.9	66.7	66.8	67.4	68.2	67.0	69.5					
7-year	39.9	44.2	43.1	43.6	44.8	49.1	54.2	54.4	54.6	58.6	60.0	65.2	65.6	65.3	66.6	65.1						
8-year	37.8	42.0	40.9	41.7	43.2	47.4	52.0	53.2	53.5	56.6	58.1	63.7	64.3	64.7	64.7							
9-year	35.7	40.1	39.1	40.1	42.2	46.4	50.4	51.3	52.3	55.2	57.1	62.8	63.1	62.9								
10-year	34.0	38.0	37.7	39.1	41.5	45.4	48.8	50.2	51.5	54.1	55.9	61.7	61.5									
11-year	32.8	36.3	36.4	38.0	40.7	43.7	47.5	49.2	50.7	52.6	54.5	60.2										
12-year	31.9	35.2	35.1	37.1	39.4	42.6	47.0	48.0	49.8	51.3	53.5											
13-year	30.7	34.2	34.2	36.1	38.8	41.9	45.4	46.5	49.0	50.0												
14-year	29.4	33.0	33.2	34.9	37.6	41.3	44.2	45.9	47.9													
15-year	28.0	31.9	32.3	34.3	36.7	40.8	43.3	45.1														
16-year	27.1	30.7	31.5	33.2	35.9	40.8	42.3															
17-year	26.5	29.9	30.8	32.5	35.6	40.3																
18-year	26.0	29.3	30.2	31.8	34.8																	
19-year	25.5	28.7	29.3	31.1																		
20-year	24.9	27.9	29.1																			

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).