

Table 19.12
Non-Hodgkin Lymphoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2013

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013
SEER 9 ^b	5,222	2,506	2,704	3,176	3,627	3,991	4,568	4,729	5,324	5,256	5,261	5,556	5,772	6,218	6,392	6,677	6,962	7,000
SEER 9 except SF-Oakland	4,285	2,088	2,267	2,603	2,819	3,089	3,505	3,694	4,158	4,246	4,363	4,509	4,817	5,201	5,392	5,606	5,813	5,956
San Francisco- Oakland SMSA ^c	937	418	437	573	808	902	1,063	1,035	1,166	1,010	898	1,047	955	1,017	1,000	1,071	1,149	1,044
San Francisco City/County	245	96	123	210	326	393	467	410	450	353	213	256	220	283	219	215	239	224

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2013

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975- 1979	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007	2008- 2009	2010- 2011	2012- 2013
SEER 9 ^b	13.4	15.5	16.1	18.3	20.1	21.5	23.3	23.3	25.2	24.4	23.8	24.4	24.6	25.7	25.4	25.4	25.4	24.3
SEER 9 except SF-Oakland	13.2	15.4	16.1	18.1	18.8	20.0	21.6	21.8	23.6	23.5	23.5	23.5	24.4	25.6	25.4	25.2	25.1	24.5
San Francisco- Oakland SMSA ^c	14.7	15.8	16.0	19.4	26.7	28.4	31.7	30.5	33.5	28.8	25.1	28.6	25.5	26.6	25.6	26.2	26.6	23.3
San Francisco City/County	14.3	14.4	17.6	29.3	44.4	51.9	61.0	52.3	57.4	44.6	27.4	32.5	26.9	35.4	26.5	25.5	27.2	25.1

^a Rates are per 100,000 males and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Non-Hodgkin Lymphoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.