

Table 19.17
Non-Hodgkin Lymphoma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.01	0.03	0.07	2.36	0.86	Asian/ Pacific Islander	0	0.01	0.03	0.06	2.11	0.84
	10	0.02	0.06	0.12	2.37	0.87		10	0.02	0.05	0.09	2.12	0.85
	20	0.04	0.10	0.23	2.36	0.87		20	0.03	0.07	0.16	2.10	0.85
	30	0.07	0.20	0.47	2.36	0.88		30	0.04	0.14	0.33	2.09	0.85
	40	0.14	0.41	0.92	2.33	0.88		40	0.10	0.29	0.68	2.06	0.85
	50	0.28	0.81	1.56	2.26	0.89		50	0.20	0.59	1.21	2.00	0.85
	60	0.57	1.37	1.99	2.13	0.90		60	0.41	1.06	1.64	1.87	0.84
	70	0.93	1.66	-	1.82	0.87		70	0.70	1.34	-	1.60	0.80
White	80	1.03	-	-	1.27	0.75	American Indian/ Alaska Native ^a	80	0.80	-	-	1.11	0.67
	0	0.01	0.03	0.06	2.46	0.90		0	0.01	0.03	0.05	1.73	0.60
	10	0.02	0.05	0.12	2.47	0.91		10	0.02	0.04	0.08	1.74	0.61
	20	0.03	0.10	0.23	2.46	0.91		20	0.02	0.06	0.14	1.73	0.61
	30	0.06	0.20	0.48	2.46	0.92		30	0.05	0.12	0.24	1.75	0.63
	40	0.14	0.42	0.96	2.43	0.92		40	0.08	0.20	0.57	1.75	0.64
	50	0.29	0.85	1.64	2.37	0.93		50	0.13	0.52	0.98	1.76	0.66
	60	0.60	1.45	2.10	2.24	0.94		60	0.43	0.93	1.51	1.79	0.69
Black	70	0.99	1.74	-	1.91	0.91	Hispanic ^b	70	0.60	1.28	-	1.61	0.69
	80	1.09	-	-	1.32	0.78		80	0.99	-	-	1.47	0.67
	0	0.01	0.03	0.07	1.38	0.49		0	0.01	0.03	0.05	2.34	0.85
	10	0.02	0.06	0.15	1.40	0.50		10	0.02	0.04	0.10	2.35	0.86
	20	0.04	0.13	0.27	1.39	0.50		20	0.03	0.08	0.20	2.35	0.86
	30	0.09	0.23	0.47	1.37	0.50		30	0.05	0.17	0.42	2.34	0.87
	40	0.14	0.39	0.72	1.32	0.50		40	0.12	0.37	0.84	2.32	0.87
	50	0.26	0.60	0.96	1.23	0.50		50	0.26	0.74	1.47	2.25	0.87
	60	0.38	0.78	1.02	1.08	0.49		60	0.51	1.28	1.92	2.11	0.87
	70	0.51	0.81	-	0.89	0.46		70	0.87	1.59	-	1.80	0.83
	80	0.49	-	-	0.61	0.39		80	0.95	-	-	1.24	0.70

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.