


Table 20.25
Cancer of the Oral Cavity and Pharynx (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013


Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.77	0.02		
High Five States				
Mississippi	6.19	0.30	01(01,03)	64.32 ^c
D.C.	5.72	0.67	02(01,22)	51.70 ^c
Louisiana	5.45	0.23	03(01,06)	44.59 ^c
Tennessee	4.86	0.18	04(02,11)	29.08 ^c
South Carolina	4.78	0.20	05(02,13)	26.82 ^c
Low Five States				
Minnesota	3.04	0.15	47(31,50)	-19.34 ^c
New Jersey	3.03	0.12	48(35,50)	-19.59 ^c
Kansas	2.97	0.20	49(29,50)	-21.24 ^c
Colorado	2.66	0.16	50(41,51)	-29.35 ^c
Utah	2.04	0.20	51(49,51)	-45.90 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	4.28	0.19	11(05,25)	13.53	Montana	3.09	0.33	45(12,51)	-17.89 ^c
Alaska	3.35	0.54	37(04,51)	-11.20	Nebraska	3.38	0.27	35(11,50)	-10.37
Arizona	3.06	0.14	46(32,50)	-18.70 ^c	Nevada	3.48	0.23	32(11,49)	-7.60
Arkansas	4.67	0.25	06(02,20)	23.91 ^c	New Hampshire	3.30	0.30	38(11,50)	-12.33
California	3.67	0.07	24(19,35)	-2.51	New Jersey	3.03	0.12	48(35,50)	-19.59 ^c
Colorado	2.66	0.16	50(41,51)	-29.35 ^c	New Mexico	3.26	0.25	41(16,50)	-13.44
Connecticut	3.15	0.18	44(24,50)	-16.45 ^c	New York	3.40	0.08	33(25,44)	-9.72
Delaware	4.32	0.43	10(02,43)	14.72	North Carolina	4.04	0.13	16(08,28)	7.16
D.C.	5.72	0.67	02(01,22)	51.70 ^c	North Dakota	3.24	0.43	42(06,51)	-13.95
Florida	4.08	0.08	12(09,23)	8.31	Ohio	3.84	0.11	19(11,34)	1.85
Georgia	4.07	0.14	15(08,29)	7.94	Oklahoma	4.07	0.21	14(06,34)	8.15
Hawaii	3.82	0.32	21(05,48)	1.36	Oregon	3.38	0.18	36(19,48)	-10.38
Idaho	3.64	0.31	27(07,49)	-3.47	Pennsylvania	3.28	0.10	40(28,47)	-12.84
Illinois	3.58	0.11	30(17,41)	-4.86	Rhode Island	3.21	0.35	43(10,50)	-14.83
Indiana	3.61	0.15	28(13,42)	-4.09	South Carolina	4.78	0.20	05(02,13)	26.82 ^c
Iowa	3.54	0.21	31(11,48)	-5.95	South Dakota	3.39	0.40	34(06,50)	-9.96
Kansas	2.97	0.20	49(29,50)	-21.24 ^c	Tennessee	4.86	0.18	04(02,11)	29.08 ^c
Kentucky	4.53	0.21	08(03,20)	20.13 ^c	Texas	4.08	0.09	13(08,23)	8.19
Louisiana	5.45	0.23	03(01,06)	44.59 ^c	Utah	2.04	0.20	51(49,51)	-45.90 ^c
Maine	4.51	0.34	09(02,32)	19.71 ^c	Vermont	3.67	0.46	23(03,50)	-2.49
Maryland	3.59	0.16	29(14,44)	-4.64	Virginia	3.65	0.14	25(13,41)	-3.11
Massachusetts	3.84	0.15	20(10,37)	1.85	Washington	3.99	0.16	17(08,32)	5.94
Michigan	3.89	0.12	18(10,33)	3.26	West Virginia	4.66	0.30	07(02,22)	23.72 ^c
Minnesota	3.04	0.15	47(31,50)	-19.34 ^c	Wisconsin	3.65	0.16	26(13,42)	-3.16
Mississippi	6.19	0.30	01(01,03)	64.32 ^c	Wyoming	3.29	0.48	39(05,51)	-12.55
Missouri	3.77	0.16	22(10,39)	-0.03					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.