

Table 21.7  
Cancer of the Ovary (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
All ages	11.9	12.5	9.6	7.5	7.8	6.5
Under 65	7.3	7.7	5.5	2.9	3.1	2.6
65 and over	43.0	45.6	37.9	39.3	40.9	33.8
All ages (IARC world std) <sup>c</sup>	8.5	8.9	6.8	4.7	4.9	4.1
<u>Age-Specific Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.2	0.2	-	-	-	-
10-14	0.8	0.7	-	-	-	-
15-19	1.5	1.5	1.4	0.1	0.1	-
20-24	1.8	1.7	1.7	0.2	0.2	-
25-29	2.4	2.5	2.2	0.3	0.3	0.3
30-34	3.3	3.3	3.2	0.5	0.6	0.4
35-39	4.7	4.9	3.2	1.1	1.1	1.1
40-44	8.8	9.2	5.5	2.2	2.2	2.0
45-49	14.4	14.9	10.0	4.6	4.9	4.0
50-54	20.0	21.0	13.8	8.5	8.9	6.9
55-59	24.3	25.8	18.8	12.9	13.5	10.8
60-64	29.7	31.5	25.2	19.2	19.9	17.0
65-69	36.9	39.3	32.1	26.6	27.7	23.6
70-74	41.7	44.4	38.9	35.2	36.6	32.5
75-79	46.8	49.7	38.7	43.7	45.6	36.7
80-84	47.8	50.9	41.2	51.7	53.5	43.5
85+	47.1	49.0	43.5	54.0	55.9	43.0

<sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.