

Table 22.9
Cancer of the Pancreas (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
1-year	14.2	15.9	17.1	18.5	17.8	19.1	20.0	19.6	22.2	21.9	23.9	25.1	22.7	23.1	26.9	25.9	27.9	29.9	29.7	30.4	31.2	34.6
2-year	5.9	6.3	6.8	8.6	8.3	9.3	9.9	8.7	10.5	10.9	11.4	11.8	11.3	11.9	12.9	13.0	13.9	15.0	15.7	16.6	15.9	
3-year	3.6	4.1	4.7	5.8	5.2	6.3	7.1	5.6	6.5	7.3	7.3	8.4	7.5	8.3	9.0	9.2	9.8	10.3	11.7	11.8		
4-year	3.0	3.3	3.8	4.8	4.0	4.8	5.5	4.5	5.6	6.0	6.1	6.7	5.7	6.4	7.2	7.9	8.3	8.4	9.7			
5-year	2.5	2.8	3.3	4.2	3.6	4.2	5.0	3.8	5.1	5.1	5.1	6.1	4.9	5.5	6.3	7.3	7.7	7.6				
6-year	2.3	2.5	2.9	4.0	3.2	3.8	4.5	3.7	4.6	4.6	4.7	5.6	4.5	5.0	5.7	6.7	7.0					
7-year	2.1	2.3	2.7	3.6	3.1	3.5	4.3	3.4	4.3	4.3	4.2	4.9	4.2	4.7	5.4	5.9						
8-year	1.9	2.1	2.5	3.4	2.7	3.4	4.1	3.1	4.0	3.8	3.9	4.7	4.0	4.3	5.0							
9-year	1.8	2.0	2.2	3.2	2.6	3.3	4.0	2.8	3.8	3.7	3.8	4.5	3.7	4.1								
10-year	1.8	1.9	2.2	3.0	2.5	3.2	3.9	2.7	3.6	3.6	3.5	4.4	3.4									
11-year	1.7	1.7	2.1	2.8	2.5	2.9	3.8	2.5	3.3	3.5	3.3	4.0										
12-year	1.7	1.7	2.1	2.7	2.3	2.8	3.6	2.4	3.2	3.1	3.1											
13-year	1.6	1.6	2.0	2.6	2.2	2.7	3.3	2.3	3.1	3.0												
14-year	1.6	1.6	1.9	2.5	1.9	2.7	3.1	1.9	3.1													
15-year	1.5	1.5	1.8	2.4	1.8	2.7	3.0	1.9														
16-year	1.5	1.4	1.7	2.2	1.7	2.2	2.9															
17-year	1.4	1.4	1.7	2.2	1.4	1.9																
18-year	1.3	1.3	1.7	2.2	1.4																	
19-year	1.3	1.3	1.6	2.1																		
20-year	1.2	1.3	1.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).