


Table 22.19  
Cancer of the Pancreas (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	12.54	0.04		
High Five States				
Louisiana	14.66	0.38	01(01,08)	16.89 <sup>c</sup>
D.C.	14.65	1.12	02(01,47)	16.88 <sup>c</sup>
Mississippi	14.34	0.47	03(01,19)	14.41 <sup>c</sup>
Maryland	13.76	0.33	04(01,24)	9.73
New Hampshire	13.50	0.64	05(01,44)	7.68
Low Five States				
New Mexico	11.19	0.47	47(27,51)	-10.73 <sup>c</sup>
Montana	11.11	0.64	48(17,51)	-11.39 <sup>c</sup>
Wyoming	11.10	0.91	49(05,51)	-11.49 <sup>c</sup>
South Dakota	10.93	0.71	50(14,51)	-12.81 <sup>c</sup>
Colorado	10.43	0.32	51(44,51)	-16.78 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	13.29	0.34	11(02,34)	6.01	Montana	11.11	0.64	48(17,51)	-11.39 <sup>c</sup>
Alaska	12.57	1.11	30(01,51)	0.25	Nebraska	12.37	0.52	35(04,49)	-1.36
Arizona	11.34	0.26	45(35,50)	-9.51	Nevada	12.17	0.45	37(07,49)	-2.93
Arkansas	13.17	0.42	12(02,39)	5.07	New Hampshire	13.50	0.64	05(01,44)	7.68
California	11.78	0.12	43(34,47)	-6.03	New Jersey	13.30	0.25	09(03,30)	6.11
Colorado	10.43	0.32	51(44,51)	-16.78 <sup>c</sup>	New Mexico	11.19	0.47	47(27,51)	-10.73 <sup>c</sup>
Connecticut	12.69	0.38	22(05,45)	1.22	New York	13.08	0.17	16(05,29)	4.36
Delaware	13.44	0.76	06(01,47)	7.21	North Carolina	12.43	0.24	34(13,43)	-0.88
D.C.	14.65	1.12	02(01,47)	16.88 <sup>c</sup>	North Dakota	12.23	0.83	36(01,51)	-2.44
Florida	11.98	0.14	40(30,46)	-4.45	Ohio	13.10	0.21	13(05,32)	4.48
Georgia	12.13	0.26	38(19,47)	-3.25	Oklahoma	12.56	0.37	31(05,46)	0.20
Hawaii	12.66	0.59	26(02,49)	0.94	Oregon	12.07	0.35	39(13,49)	-3.77
Idaho	12.80	0.59	21(02,48)	2.08	Pennsylvania	13.30	0.20	10(03,25)	6.04
Illinois	12.89	0.21	19(07,36)	2.84	Rhode Island	12.89	0.70	20(01,49)	2.80
Indiana	13.06	0.30	17(04,37)	4.18	South Carolina	13.09	0.35	14(03,39)	4.44
Iowa	12.53	0.39	32(05,46)	-0.04	South Dakota	10.93	0.71	50(14,51)	-12.81 <sup>c</sup>
Kansas	12.60	0.43	29(04,47)	0.46	Tennessee	12.69	0.29	24(06,41)	1.20
Kentucky	13.03	0.36	18(03,39)	3.90	Texas	11.69	0.16	44(34,49)	-6.74
Louisiana	14.66	0.38	01(01,08)	16.89 <sup>c</sup>	Utah	11.27	0.48	46(22,51)	-10.12 <sup>c</sup>
Maine	11.84	0.56	42(07,51)	-5.60	Vermont	13.36	0.90	08(01,49)	6.56
Maryland	13.76	0.33	04(01,24)	9.73	Virginia	12.65	0.27	27(07,41)	0.90
Massachusetts	12.67	0.28	25(07,42)	1.08	Washington	12.63	0.29	28(07,42)	0.70
Michigan	13.44	0.23	07(03,25)	7.17	West Virginia	11.97	0.48	41(09,50)	-4.52
Minnesota	12.43	0.31	33(09,46)	-0.84	Wisconsin	13.09	0.30	15(03,36)	4.44
Mississippi	14.34	0.47	03(01,19)	14.41 <sup>c</sup>	Wyoming	11.10	0.91	49(05,51)	-11.49 <sup>c</sup>
Missouri	12.69	0.29	23(06,41)	1.21					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.