

Table 23.7  
Cancer of the Prostate (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
All ages	129.4	121.9	203.5	20.7	19.1	44.2
Under 65	51.4	47.7	93.0	1.7	1.4	4.2
65 and over	668.5	635.3	967.5	151.7	141.3	320.9
All ages (IARC world std) <sup>c</sup>	90.2	84.9	145.8	9.0	8.2	20.1
<u>Age-Specific Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	1.0	0.8	2.9	0.0	-	-
40-44	9.8	8.5	23.2	0.2	0.1	0.4
45-49	42.5	35.9	102.5	0.9	0.8	2.0
50-54	132.4	121.2	250.5	3.2	2.7	8.0
55-59	285.0	264.9	506.3	8.7	7.4	21.6
60-64	486.3	459.8	810.7	19.7	17.0	49.2
65-69	746.8	710.9	1,153.1	39.3	34.4	99.7
70-74	782.8	746.2	1,113.4	74.9	66.9	182.2
75-79	686.3	648.8	942.1	133.5	122.5	295.7
80-84	504.7	478.2	684.2	234.0	219.2	501.7
85+	418.4	398.0	628.4	494.6	473.5	929.5

<sup>a</sup> SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.