

Table 23.8  
Cancer of the Prostate (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Diagnosis Year, Stage and Age

	All Races, Males			White Males			Black Males		
	All	<65	65+	All	<65	65+	All	<65	65+
<b>5-Year Relative Survival (Percent)</b>									
<b>Year of Diagnosis:</b>									
1960-1963 <sup>a</sup>	-	-	-	50	-	-	35	-	-
1970-1973 <sup>a</sup>	-	-	-	63	-	-	55	-	-
1975-1977 <sup>b</sup>	67.8	73.4	66.1	68.5	75.2	66.6	60.7	62.0	60.1
1978-1980 <sup>b</sup>	70.1	75.2	68.6	71.2	76.7	69.6	62.1	65.0	60.6
1981-1983 <sup>b</sup>	72.0	76.0	70.9	73.1	77.6	71.9	62.8	65.0	61.9
1984-1986 <sup>b</sup>	75.2	77.3	74.6	76.4	79.0	75.7	65.6	67.2	64.9
1987-1989 <sup>b</sup>	82.8	83.2	82.7	84.4	84.7	84.3	71.1	72.8	70.6
1990-1992 <sup>b</sup>	92.9	93.0	92.9	94.1	94.0	94.1	84.4	86.8	83.5
1993-1995 <sup>b</sup>	95.2	95.9	94.9	95.9	96.6	95.6	91.7	93.3	90.8
1996-1998 <sup>b</sup>	97.4	98.5	96.7	97.8	98.9	97.2	94.9	96.7	93.4
1999-2001 <sup>b</sup>	99.3	99.1	99.3	99.7	99.5	99.6	97.3	98.0	96.2
2002-2005 <sup>b</sup>	99.6	99.4	99.7	99.8	99.6	99.7	97.7	98.8	96.3
2006-2012 <sup>b</sup>	99.3 <sup>f</sup>	99.5 <sup>f</sup>	99.2 <sup>f</sup>	99.7 <sup>f</sup>	99.8 <sup>f</sup>	99.4 <sup>f</sup>	97.3 <sup>e</sup>	98.1 <sup>f</sup>	96.1 <sup>f</sup>
<b>5-Year Period Survival (Percent)<sup>cd</sup></b>									
2012	98.7	99.1	98.4	98.9	99.3	98.5	97.1	98.1	95.7
<b>Stage Distribution (%) 2006-2012<sup>ce</sup></b>									
<b>All Stages</b>									
Number of cases	368,301	163,823	204,478	282,870	122,545	160,325	54,632	30,066	24,566
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	80	80	80	80	79	81	82	81	82
Regional	12	16	9	13	16	10	10	13	7
Distant	4	3	6	4	3	5	5	4	7
Unstaged	4	2	5	3	1	4	3	2	4
<b>5-Year Relative Survival (Percent), 2006-2012<sup>c</sup></b>									
<b>Age at Diagnosis:</b>									
Ages <45	96.1	-	-	95.4	-	-	97.4	-	-
Ages 45-54	98.1	-	-	98.2	-	-	97.5	-	-
Ages 55-64	99.4	-	-	99.7	-	-	98.2	-	-
Ages 65-74	100.0	-	-	100.0	-	-	97.5	-	-
Ages 75+	95.1	-	-	95.0	-	-	90.5	-	-
Ages <65	99.1	-	-	99.3	-	-	98.0	-	-
Ages 65+	98.7	-	-	99.0	-	-	95.6	-	-
<b>Stage<sup>e</sup>:</b>									
All Stages	98.9	99.1	98.7	99.2	99.3	99.0	96.9	98.0	95.6
Localized	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Regional	100.0	100.0	100.0	100.0	99.9	100.0	100.0	100.0	98.7
Distant	29.3	32.5	27.9	28.7	31.6	27.6	29.0	32.4	26.3
Unstaged	79.4	89.7	76.3	74.4	87.4	70.8	68.0	82.2	61.8

<sup>a</sup> Based on End Results data from a series of hospital registries and one population-based registry.

<sup>b</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

<sup>c</sup> Based on follow-up of patients into 2013.

<sup>d</sup> SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

<sup>e</sup> Based on follow-up of patients into 2013.

<sup>f</sup> Period survival provides a 2012 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival), (2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival).

<sup>g</sup> Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

<sup>h</sup> The difference between 1975-1977 and 2006-2012 is statistically significant ( $p < .05$ ).

<sup>i</sup> The standard error is between 5 and 10 percentage points.

<sup>j</sup> The standard error is greater than 10 percentage points.

<sup>k</sup> Statistic could not be calculated due to fewer than 25 cases during the time period.