


Table 23.14
Cancer of the Prostate (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013


Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	20.65	0.06		
High Five States				
D.C.	35.66	1.84	01(01,01)	72.66 ^c
Mississippi	27.37	0.70	02(02,03)	32.53 ^c
Alabama	25.17	0.51	03(02,06)	21.87 ^c
South Carolina	24.27	0.52	04(03,09)	17.53 ^c
Georgia	23.87	0.40	05(03,09)	15.58 ^c
Low Five States				
Kansas	18.92	0.54	47(31,50)	-8.40
Wyoming	18.91	1.27	48(09,50)	-8.42
Arizona	18.69	0.35	49(37,50)	-9.50
Florida	18.11	0.18	50(46,50)	-12.32 ^c
Hawaii	13.94	0.63	51(51,51)	-32.52 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	25.17	0.51	03(02,06)	21.87 ^c	Montana	22.51	0.94	08(03,38)	8.99
Alaska	20.48	1.62	32(03,50)	-0.84	Nebraska	21.62	0.71	17(06,41)	4.68
Arizona	18.69	0.35	49(37,50)	-9.50	Nevada	21.29	0.65	23(07,44)	3.11
Arkansas	21.80	0.58	15(06,36)	5.57	New Hampshire	20.05	0.83	36(09,50)	-2.94
California	20.55	0.17	29(24,37)	-0.48	New Jersey	19.47	0.32	43(32,49)	-5.74
Colorado	22.03	0.50	14(06,32)	6.68	New Mexico	21.42	0.69	21(06,42)	3.72
Connecticut	19.22	0.47	44(30,50)	-6.96	New York	19.93	0.22	38(30,44)	-3.50
Delaware	20.05	0.99	35(07,50)	-2.91	North Carolina	22.21	0.35	11(06,25)	7.53
D.C.	35.66	1.84	01(01,01)	72.66 ^c	North Dakota	19.97	1.08	37(07,50)	-3.30
Florida	18.11	0.18	50(46,50)	-12.32 ^c	Ohio	20.92	0.28	28(16,37)	1.30
Georgia	23.87	0.40	05(03,09)	15.58 ^c	Oklahoma	21.63	0.52	16(07,35)	4.72
Hawaii	13.94	0.63	51(51,51)	-32.52 ^c	Oregon	22.05	0.49	13(06,31)	6.75
Idaho	23.65	0.85	06(03,25)	14.49 ^c	Pennsylvania	20.38	0.25	33(23,41)	-1.29
Illinois	21.52	0.29	19(10,31)	4.19	Rhode Island	20.54	0.90	30(07,50)	-0.55
Indiana	21.18	0.40	26(10,37)	2.58	South Carolina	24.27	0.52	04(03,09)	17.53 ^c
Iowa	19.82	0.51	40(22,49)	-4.04	South Dakota	20.27	0.99	34(07,50)	-1.87
Kansas	18.92	0.54	47(31,50)	-8.40	Tennessee	21.47	0.41	20(08,35)	3.96
Kentucky	20.52	0.49	31(14,45)	-0.66	Texas	19.19	0.22	45(36,49)	-7.08
Louisiana	23.14	0.52	07(04,20)	12.07 ^c	Utah	22.32	0.72	10(04,35)	8.06
Maine	21.10	0.78	27(06,47)	2.17	Vermont	21.27	1.19	25(03,50)	3.01
Maryland	21.29	0.43	24(09,36)	3.07	Virginia	22.19	0.38	12(06,27)	7.46
Massachusetts	19.88	0.36	39(26,48)	-3.73	Washington	21.33	0.39	22(09,36)	3.28
Michigan	19.68	0.30	41(30,47)	-4.73	West Virginia	19.56	0.65	42(20,50)	-5.29
Minnesota	21.56	0.43	18(08,35)	4.40	Wisconsin	22.41	0.41	09(06,25)	8.51
Mississippi	27.37	0.70	02(02,03)	32.53 ^c	Wyoming	18.91	1.27	48(09,50)	-8.42
Missouri	18.97	0.38	46(35,50)	-8.12					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.