

Table 25.4  
Cancer of the Testis (Invasive)

Delay-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Age

|                    | All Races, Males |      |      | White Males |      |      | Black Males |      |      |
|--------------------|------------------|------|------|-------------|------|------|-------------|------|------|
|                    | All              | <50  | 50+  | All         | <50  | 50+  | All         | <50  | 50+  |
| Year of Diagnosis: |                  |      |      |             |      |      |             |      |      |
| 1975               | 3.73             | 4.23 | 2.41 | 4.16        | 4.74 | 2.65 | 0.68        | 0.74 | 0.50 |
| 1976               | 3.44             | 4.18 | 1.51 | 3.77        | 4.56 | 1.72 | 1.15        | 1.60 | 0.00 |
| 1977               | 4.30             | 5.09 | 2.24 | 4.88        | 5.82 | 2.41 | 0.48        | 0.46 | 0.55 |
| 1978               | 3.58             | 4.24 | 1.84 | 3.95        | 4.65 | 2.11 | 1.49        | 2.06 | 0.00 |
| 1979               | 3.88             | 4.54 | 2.15 | 4.32        | 5.09 | 2.31 | 0.78        | 0.71 | 0.96 |
| 1980               | 4.35             | 5.28 | 1.89 | 4.90        | 5.96 | 2.11 | 0.83        | 1.15 | 0.00 |
| 1981               | 4.23             | 5.13 | 1.85 | 4.77        | 5.79 | 2.07 | 0.60        | 0.83 | 0.00 |
| 1982               | 4.40             | 5.23 | 2.25 | 5.01        | 5.99 | 2.44 | 0.70        | 0.80 | 0.45 |
| 1983               | 4.60             | 5.50 | 2.24 | 5.23        | 6.30 | 2.43 | 1.29        | 1.04 | 1.97 |
| 1984               | 4.40             | 5.36 | 1.86 | 5.04        | 6.17 | 2.08 | 0.62        | 0.85 | 0.00 |
| 1985               | 4.47             | 5.47 | 1.83 | 5.13        | 6.31 | 2.05 | 1.07        | 1.32 | 0.42 |
| 1986               | 4.82             | 5.83 | 2.17 | 5.57        | 6.79 | 2.39 | 0.56        | 0.56 | 0.55 |
| 1987               | 5.05             | 6.27 | 1.86 | 5.70        | 7.13 | 1.94 | 1.08        | 1.32 | 0.44 |
| 1988               | 4.63             | 5.83 | 1.46 | 5.44        | 6.86 | 1.72 | 0.60        | 0.82 | 0.00 |
| 1989               | 5.48             | 6.71 | 2.24 | 6.31        | 7.77 | 2.46 | 1.23        | 1.13 | 1.46 |
| 1990               | 5.09             | 6.27 | 2.02 | 5.99        | 7.40 | 2.30 | 0.56        | 0.59 | 0.47 |
| 1991               | 5.10             | 6.24 | 2.12 | 5.75        | 7.07 | 2.31 | 1.18        | 1.17 | 1.19 |
| 1992               | 5.15             | 6.44 | 1.76 | 6.09        | 7.66 | 1.97 | 0.53        | 0.73 | 0.00 |
| 1993               | 5.07             | 6.37 | 1.65 | 5.99        | 7.54 | 1.93 | 0.66        | 0.91 | 0.00 |
| 1994               | 5.48             | 6.76 | 2.12 | 6.29        | 7.81 | 2.30 | 1.23        | 1.54 | 0.42 |
| 1995               | 4.55             | 5.57 | 1.88 | 5.23        | 6.42 | 2.11 | 1.46        | 1.87 | 0.40 |
| 1996               | 5.25             | 6.60 | 1.73 | 6.20        | 7.85 | 1.89 | 1.10        | 1.05 | 1.21 |
| 1997               | 5.40             | 6.52 | 2.46 | 6.34        | 7.73 | 2.68 | 1.27        | 1.24 | 1.34 |
| 1998               | 5.60             | 7.12 | 1.62 | 6.58        | 8.41 | 1.79 | 1.57        | 2.05 | 0.33 |
| 1999               | 5.45             | 6.84 | 1.81 | 6.34        | 7.99 | 2.02 | 1.71        | 2.02 | 0.92 |
| 2000               | 5.74             | 7.17 | 1.99 | 6.74        | 8.42 | 2.33 | 1.52        | 1.86 | 0.62 |
| 2001               | 5.53             | 6.86 | 2.04 | 6.52        | 8.15 | 2.24 | 1.79        | 1.99 | 1.29 |
| 2002               | 5.81             | 7.27 | 1.98 | 6.95        | 8.70 | 2.34 | 1.39        | 1.80 | 0.29 |
| 2003               | 5.35             | 6.69 | 1.83 | 6.49        | 8.21 | 1.98 | 1.14        | 1.27 | 0.80 |
| 2004               | 5.97             | 7.38 | 2.27 | 7.35        | 9.15 | 2.63 | 1.05        | 1.21 | 0.63 |
| 2005               | 6.03             | 7.52 | 2.13 | 7.29        | 9.19 | 2.30 | 1.02        | 1.32 | 0.24 |
| 2006               | 5.67             | 7.03 | 2.11 | 6.79        | 8.44 | 2.45 | 1.62        | 1.72 | 1.37 |
| 2007               | 6.04             | 7.45 | 2.33 | 7.24        | 8.99 | 2.65 | 1.48        | 1.57 | 1.26 |
| 2008               | 6.10             | 7.39 | 2.74 | 7.41        | 9.09 | 3.03 | 1.45        | 1.75 | 0.67 |
| 2009               | 6.11             | 7.55 | 2.34 | 7.33        | 9.18 | 2.48 | 1.70        | 1.72 | 1.66 |
| 2010               | 5.88             | 7.17 | 2.51 | 6.92        | 8.47 | 2.85 | 1.71        | 1.94 | 1.09 |
| 2011               | 5.95             | 7.45 | 2.03 | 7.17        | 9.05 | 2.26 | 1.49        | 1.99 | 0.20 |
| 2012               | 6.02             | 7.33 | 2.59 | 7.33        | 9.00 | 2.95 | 1.29        | 1.41 | 0.97 |
| 2013               | 6.14             | 7.54 | 2.47 | 7.51        | 9.27 | 2.91 | 1.65        | 1.97 | 0.81 |

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

- <sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).  
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
 - Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.