

Table 25.5
Cancer of the Testis (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Age

	All Races, Males			White Males			Black Males		
	All	<50	50+	All	<50	50+	All	<50	50+
Year of Diagnosis:									
1975-2013	5.22	6.42	2.10	6.08	7.51	2.34	1.22	1.41	0.72
1975	3.73	4.23	2.41	4.16	4.74	2.65	-	-	-
1976	3.44	4.18	1.51	3.77	4.56	1.72	-	-	-
1977	4.30	5.09	2.24	4.88	5.82	2.41	-	-	-
1978	3.58	4.24	1.84	3.95	4.65	2.11	-	-	-
1979	3.88	4.54	2.15	4.32	5.09	2.31	-	-	-
1980	4.35	5.28	1.89	4.90	5.96	2.11	-	-	-
1981	4.23	5.13	1.85	4.77	5.79	2.07	-	-	-
1982	4.40	5.23	2.25	5.01	5.99	2.44	-	-	-
1983	4.60	5.50	2.24	5.23	6.30	2.43	-	-	-
1984	4.40	5.36	1.86	5.04	6.17	2.08	-	-	-
1985	4.47	5.47	1.83	5.13	6.31	2.05	-	-	-
1986	4.82	5.83	2.17	5.57	6.79	2.39	-	-	-
1987	5.05	6.27	1.86	5.70	7.13	1.94	-	-	-
1988	4.63	5.83	1.46	5.44	6.86	1.72	-	-	-
1989	5.48	6.71	2.24	6.31	7.77	2.46	-	-	-
1990	5.09	6.27	2.02	5.99	7.40	2.30	-	-	-
1991	5.10	6.24	2.12	5.75	7.07	2.31	-	-	-
1992	5.15	6.44	1.76	6.09	7.66	1.97	-	-	-
1993	5.07	6.37	1.65	5.99	7.54	1.93	-	-	-
1994	5.48	6.76	2.12	6.29	7.81	2.30	1.23	1.54	-
1995	4.55	5.57	1.88	5.23	6.42	2.11	1.46	1.87	-
1996	5.25	6.60	1.73	6.20	7.85	1.89	-	-	-
1997	5.40	6.52	2.46	6.34	7.73	2.68	1.27	-	-
1998	5.60	7.12	1.62	6.58	8.41	1.79	1.57	2.05	-
1999	5.45	6.84	1.81	6.34	7.99	2.02	1.71	2.02	-
2000	5.74	7.17	1.99	6.74	8.42	2.33	1.52	1.86	-
2001	5.53	6.86	2.04	6.52	8.15	2.24	1.79	1.99	-
2002	5.81	7.27	1.98	6.95	8.70	2.34	1.39	1.80	-
2003	5.35	6.69	1.83	6.49	8.21	1.98	1.14	-	-
2004	5.97	7.38	2.27	7.35	9.15	2.63	1.05	-	-
2005	6.03	7.52	2.13	7.28	9.18	2.30	1.02	1.32	-
2006	5.66	7.02	2.10	6.78	8.43	2.45	1.62	1.71	-
2007	6.03	7.44	2.32	7.22	8.97	2.64	1.48	1.56	-
2008	6.09	7.37	2.73	7.39	9.06	3.02	1.45	1.74	-
2009	6.09	7.53	2.33	7.30	9.14	2.47	1.70	1.71	-
2010	5.86	7.15	2.50	6.89	8.44	2.84	1.70	1.93	-
2011	5.92	7.41	2.02	7.13	9.00	2.25	1.48	1.97	-
2012	5.97	7.27	2.57	7.27	8.93	2.92	1.28	1.40	-
2013	6.05	7.43	2.43	7.37	9.10	2.85	1.61	1.93	-

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.