

Table 25.7
Cancer of the Testis (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
All ages	5.7	6.7	1.5	0.3	0.3	0.1
Under 65	6.3	7.5	1.6	0.2	0.3	0.1
65 and over	1.2	1.3	-	0.3	0.3	-
All ages (IARC world std) ^c	5.4	6.4	1.3	0.2	0.2	0.1
<u>Age-Specific Rates, 2009-2013</u>						
Age at Diagnosis/Death:						
<1	0.9	0.9	-	-	-	-
1-4	0.5	0.5	-	-	-	-
5-9	-	-	-	-	-	-
10-14	0.2	0.2	-	-	-	-
15-19	4.0	5.1	-	0.1	0.1	-
20-24	10.5	12.9	1.9	0.3	0.4	-
25-29	14.9	17.9	2.6	0.5	0.6	-
30-34	13.8	16.6	3.4	0.4	0.5	-
35-39	11.0	13.2	3.1	0.4	0.4	-
40-44	8.1	9.4	2.9	0.3	0.4	-
45-49	6.6	7.8	1.8	0.4	0.5	-
50-54	4.0	4.6	1.6	0.3	0.4	-
55-59	3.2	3.6	1.5	0.3	0.3	-
60-64	2.1	2.5	-	0.2	0.2	-
65-69	1.3	1.4	-	0.2	0.2	-
70-74	1.1	1.3	-	0.2	0.2	-
75-79	1.2	1.3	-	0.3	0.3	-
80-84	0.9	1.1	-	0.4	0.4	-
85+	1.0	1.0	-	0.6	0.6	-

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.