

Table 25.10
Cancer of the Testis (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.03	0.14	0.39	0.02	Asian/ Pacific Islander	0	0.00	0.01	0.06	0.15	0.01
	10	0.02	0.14	0.26	0.39	0.02		10	0.01	0.06	0.10	0.15	0.01
	20	0.12	0.24	0.31	0.37	0.02		20	0.05	0.09	0.12	0.14	0.01
	30	0.12	0.19	0.22	0.25	0.02		30	0.04	0.07	0.08	0.09	0.01
	40	0.07	0.10	0.12	0.13	0.01		40	0.02	0.04	0.04	0.04	0.00
	50	0.03	0.05	0.06	0.06	0.01		50	0.01	0.02	0.02	0.02	0.00
	60	0.02	0.02	0.03	0.03	0.01		60	0.01	0.01	0.01	0.01	0.00
	70	0.01	0.01	-	0.01	0.01		70	0.00	0.00	-	0.00	0.00
White	80	0.01	-	-	0.01	0.00	American Indian/ Alaska Native ^a	80	0.00	-	-	0.00	0.00
	0	0.00	0.03	0.18	0.46	0.02		0	0.00	0.02	0.17	0.35	0.02
	10	0.03	0.17	0.32	0.46	0.02		10	0.02	0.16	0.26	0.35	0.02
	20	0.14	0.29	0.37	0.43	0.02		20	0.15	0.25	0.30	0.33	0.02
	30	0.14	0.23	0.26	0.29	0.02		30	0.10	0.15	0.17	0.19	0.01
	40	0.08	0.12	0.14	0.15	0.01		40	0.06	0.08	0.09	0.09	0.01
	50	0.04	0.06	0.07	0.07	0.01		50	0.02	0.04	0.04	0.04	0.01
	60	0.02	0.03	0.03	0.03	0.01		60	0.02	0.02	0.02	0.02	0.01
Black	70	0.01	0.01	-	0.02	0.01	Hispanic ^b	70	0.00	0.00	-	0.00	0.00
	80	0.01	-	-	0.01	0.00		80	0.00	-	-	0.00	0.00
	0	0.00	0.01	0.03	0.10	0.01		0	0.00	0.04	0.17	0.35	0.02
	10	0.00	0.02	0.06	0.10	0.01		10	0.03	0.17	0.26	0.35	0.02
	20	0.02	0.05	0.08	0.10	0.01		20	0.13	0.23	0.27	0.32	0.02
	30	0.03	0.06	0.07	0.08	0.01		30	0.10	0.14	0.16	0.19	0.01
	40	0.02	0.04	0.04	0.05	0.01		40	0.04	0.06	0.08	0.09	0.01
	50	0.02	0.02	0.02	0.03	0.01		50	0.02	0.03	0.05	0.05	0.01
	60	0.01	0.01	0.01	0.01	0.00		60	0.01	0.03	0.03	0.03	0.01
	70	0.00	0.01	-	0.01	0.00		70	0.02	0.02	-	0.02	0.00
	80	0.00	-	-	0.00	0.00		80	0.00	-	-	0.01	0.00

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.