



Table 25.14  
Cancer of the Testis (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	0.25	0.01		
High Five States				
West Virginia	0.39	0.10	01(01,30)	55.81 <sup>c</sup>
Oklahoma	0.39	0.07	02(01,22)	54.20 <sup>c</sup>
Arizona	0.38	0.05	03(01,15)	51.90 <sup>c</sup>
Missouri	0.36	0.05	04(01,21)	43.18 <sup>c</sup>
California	0.32	0.02	05(02,14)	27.25 <sup>c</sup>
Low Five States				
South Carolina	0.19	0.04	28(07,32)	-24.23 <sup>c</sup>
Maryland	0.19	0.04	29(07,32)	-24.54 <sup>c</sup>
Massachusetts	0.15	0.03	30(17,32)	-41.15 <sup>c</sup>
Virginia	0.15	0.03	31(20,32)	-42.08 <sup>c</sup>
Minnesota	0.13	0.03	32(20,32)	-47.33 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	0.24	0.05	18(02,32)	-4.24	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.38	0.05	03(01,15)	51.90 <sup>c</sup>	Nevada	0.28	0.06	10(01,31)	12.39
Arkansas	-	-	-(-,-)	-	New Hampshire	-	-	-(-,-)	-
California	0.32	0.02	05(02,14)	27.25 <sup>c</sup>	New Jersey	0.23	0.03	20(05,31)	-7.18
Colorado	0.29	0.05	09(01,29)	14.39	New Mexico	0.31	0.08	07(01,32)	22.98 <sup>c</sup>
Connecticut	-	-	-(-,-)	-	New York	0.22	0.02	23(10,30)	-13.33
Delaware	-	-	-(-,-)	-	North Carolina	0.25	0.03	16(04,29)	-0.32
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.27	0.02	12(04,25)	7.28	Ohio	0.25	0.03	17(04,29)	-0.74
Georgia	0.21	0.03	26(08,32)	-16.74 <sup>c</sup>	Oklahoma	0.39	0.07	02(01,22)	54.20 <sup>c</sup>
Hawaii	-	-	-(-,-)	-	Oregon	-	-	-(-,-)	-
Idaho	-	-	-(-,-)	-	Pennsylvania	0.26	0.03	15(04,28)	2.00
Illinois	0.19	0.02	27(13,32)	-23.79 <sup>c</sup>	Rhode Island	-	-	-(-,-)	-
Indiana	0.30	0.05	08(01,27)	21.04 <sup>c</sup>	South Carolina	0.19	0.04	28(07,32)	-24.23 <sup>c</sup>
Iowa	0.26	0.06	13(01,32)	4.72	South Dakota	-	-	-(-,-)	-
Kansas	0.22	0.06	24(02,32)	-13.36	Tennessee	0.23	0.04	19(03,31)	-6.63
Kentucky	0.31	0.05	06(01,28)	23.24 <sup>c</sup>	Texas	0.28	0.02	11(04,23)	10.17
Louisiana	0.21	0.04	25(04,32)	-16.58 <sup>c</sup>	Utah	-	-	-(-,-)	-
Maine	-	-	-(-,-)	-	Vermont	-	-	-(-,-)	-
Maryland	0.19	0.04	29(07,32)	-24.54 <sup>c</sup>	Virginia	0.15	0.03	31(20,32)	-42.08 <sup>c</sup>
Massachusetts	0.15	0.03	30(17,32)	-41.15 <sup>c</sup>	Washington	0.22	0.04	22(05,32)	-11.48
Michigan	0.26	0.03	14(03,29)	2.54	West Virginia	0.39	0.10	01(01,30)	55.81 <sup>c</sup>
Minnesota	0.13	0.03	32(20,32)	-47.33 <sup>c</sup>	Wisconsin	0.23	0.04	21(03,32)	-7.98
Mississippi	-	-	-(-,-)	-	Wyoming	-	-	-(-,-)	-
Missouri	0.36	0.05	04(01,21)	43.18 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.