

Table 25.15
Cancer of the Testis (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2013
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <21 ^e	0 to <38 ^e	>=38 ^g	Complete ^h
<u>Race</u>										
All Races ^b	42,726	39,615	37,513	32,672	28,999	22,670	170,263	226,015	14,357	240,372
White ^b	39,092	37,146	35,108	31,004	27,916	21,895	159,376	213,123	13,723	226,846
Black ^b	1,303	971	1,089	663	368	298	4,245	5,135	150	5,285
Asian/Pacific Islander ^c	936	708	648	+	+	+	2,868	+	+	+
Hispanic ^d	6,848	5,262	3,793	+	+	+	19,147	+	+	+

Estimated prevalence percent^a on January 1, 2013, of the SEER^c population diagnosed in the previous 21 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0966%	0.0025%	0.0070%	0.0709%	0.1733%	0.2172%	0.1591%	0.0754%	0.0291%	0.0148%	0.0961%
White ^c	0.1192%	0.0024%	0.0083%	0.0884%	0.2171%	0.2692%	0.1921%	0.0900%	0.0345%	0.0174%	0.1185%
Black ^c	0.0209%	0.0019%	0.0023%	0.0140%	0.0312%	0.0508%	0.0424%	0.0206%	-	-	0.0216%
Asian/Pacific Islander ^c	0.0324%	0.0040%	0.0047%	0.0298%	0.0519%	0.0646%	0.0518%	0.0235%	0.0087%	-	0.0311%
Hispanic ^d	0.0717%	0.0029%	0.0089%	0.0785%	0.1583%	0.1462%	0.0931%	0.0364%	0.0183%	-	0.0728%

^a US 2013 cancer prevalence counts are based on 2013 cancer prevalence proportions from the SEER registries and 1/1/2013 US population estimates based on the average of 2012 and 2013 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 13 Areas excluding the Alaska Native Registry

(d) NHIA for Hispanic for SEER 13 Areas excluding the Alaska Native Registry.

^e Maximum limited-duration prevalence: 38 years for 1975-2013 SEER 9 data; 21 years for 1992-2013 SEER 13 data (excluding the Alaska Native Registry) used to calculate prevalence for Hispanics and Asian Pacific Islanders.

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 38 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <38 and >=38. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.