

Table 26.13
Cancer of the Thyroid (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2011-2013 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.01	0.03	0.62	0.06	Asian/ Pacific Islander	0	0.00	0.00	0.03	0.67	0.09
	10	0.01	0.03	0.10	0.62	0.06		10	0.00	0.03	0.09	0.67	0.10
	20	0.03	0.09	0.18	0.62	0.06		20	0.02	0.09	0.19	0.67	0.10
	30	0.06	0.16	0.28	0.60	0.06		30	0.07	0.16	0.28	0.65	0.10
	40	0.10	0.22	0.36	0.54	0.06		40	0.10	0.21	0.35	0.59	0.10
	50	0.13	0.28	0.40	0.46	0.06		50	0.12	0.26	0.40	0.50	0.10
	60	0.16	0.30	0.36	0.36	0.06		60	0.15	0.30	0.38	0.40	0.09
	70	0.16	0.23	-	0.23	0.05		70	0.16	0.25	-	0.27	0.08
White	80	0.09	-	-	0.10	0.04	American Indian/ Alaska Native ^a	80	0.11	-	-	0.13	0.06
	0	0.00	0.01	0.04	0.65	0.06		0	0.00	0.00	0.01	0.34	0.07
	10	0.01	0.04	0.11	0.66	0.06		10	0.00	0.01	0.05	0.34	0.07
	20	0.03	0.10	0.20	0.65	0.06		20	0.01	0.05	0.11	0.34	0.07
	30	0.07	0.17	0.30	0.63	0.06		30	0.05	0.10	0.19	0.35	0.07
	40	0.10	0.23	0.39	0.57	0.06		40	0.06	0.15	0.23	0.31	0.07
	50	0.13	0.29	0.42	0.48	0.06		50	0.10	0.18	0.26	0.26	0.08
	60	0.17	0.31	0.37	0.38	0.06		60	0.09	0.18	0.18	0.18	0.08
Black	70	0.17	0.23	-	0.24	0.05	Hispanic ^b	70	0.10	0.10	-	0.10	0.08
	80	0.10	-	-	0.11	0.04		80	0.01	-	-	0.01	0.07
	0	0.00	0.00	0.01	0.29	0.03		0	0.00	0.01	0.02	0.49	0.06
	10	0.00	0.01	0.03	0.29	0.03		10	0.01	0.02	0.06	0.49	0.06
	20	0.01	0.03	0.07	0.29	0.03		20	0.02	0.06	0.12	0.49	0.06
	30	0.02	0.07	0.12	0.29	0.03		30	0.04	0.10	0.19	0.48	0.06
	40	0.05	0.10	0.19	0.28	0.03		40	0.06	0.15	0.27	0.44	0.07
	50	0.06	0.15	0.22	0.24	0.03		50	0.09	0.21	0.32	0.39	0.07
	60	0.10	0.18	0.20	0.20	0.03		60	0.13	0.24	0.31	0.31	0.06
	70	0.10	0.13	-	0.14	0.03		70	0.13	0.20	-	0.21	0.05
	80	0.05	-	-	0.05	0.02		80	0.10	-	-	0.11	0.04

Devcan 6.7.4, August 2016, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.