


Table 26.18
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013


Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
Hawaii	0.79	0.10	01(01,23)	55.15 ^c
D.C.	0.76	0.16	02(01,50)	49.62 ^c
Idaho	0.68	0.09	03(01,45)	33.71 ^c
Alaska	0.66	0.17	04(01,50)	29.24 ^c
Rhode Island	0.63	0.10	05(01,49)	24.42 ^c
Low Five States				
Mississippi	0.43	0.05	46(07,50)	-15.90 ^c
Maryland	0.43	0.04	47(14,50)	-15.97 ^c
Florida	0.43	0.02	48(29,50)	-16.10 ^c
Connecticut	0.42	0.05	49(10,50)	-16.62 ^c
Georgia	0.41	0.03	50(25,50)	-20.01 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.45	0.04	39(08,50)	-10.44	Montana	0.47	0.09	35(02,50)	-7.97
Alaska	0.66	0.17	04(01,50)	29.24 ^c	Nebraska	0.57	0.07	08(01,48)	13.20
Arizona	0.47	0.04	34(08,50)	-7.90	Nevada	0.53	0.06	14(02,49)	5.27
Arkansas	0.46	0.05	38(05,50)	-10.07	New Hampshire	0.45	0.08	41(03,50)	-10.54
California	0.59	0.02	06(03,18)	16.23 ^c	New Jersey	0.50	0.03	27(06,46)	-2.29
Colorado	0.49	0.05	30(05,49)	-3.03	New Mexico	0.57	0.07	10(01,49)	11.80
Connecticut	0.42	0.05	49(10,50)	-16.62 ^c	New York	0.55	0.02	12(05,30)	8.51
Delaware	0.58	0.11	07(01,50)	13.98	North Carolina	0.51	0.03	18(05,44)	0.78
D.C.	0.76	0.16	02(01,50)	49.62 ^c	North Dakota	0.46	0.10	37(01,50)	-8.38
Florida	0.43	0.02	48(29,50)	-16.10 ^c	Ohio	0.52	0.03	15(06,40)	2.70
Georgia	0.41	0.03	50(25,50)	-20.01 ^c	Oklahoma	0.47	0.05	36(05,50)	-8.11
Hawaii	0.79	0.10	01(01,23)	55.15 ^c	Oregon	0.52	0.05	16(03,49)	2.29
Idaho	0.68	0.09	03(01,45)	33.71 ^c	Pennsylvania	0.52	0.03	17(07,40)	1.53
Illinois	0.51	0.03	22(07,44)	-0.32	Rhode Island	0.63	0.10	05(01,49)	24.42 ^c
Indiana	0.47	0.04	31(08,49)	-6.71	South Carolina	0.43	0.04	44(10,50)	-14.69
Iowa	0.47	0.05	32(05,50)	-7.72	South Dakota	0.47	0.10	33(01,50)	-7.86
Kansas	0.50	0.06	25(03,50)	-1.20	Tennessee	0.44	0.04	42(10,50)	-12.52
Kentucky	0.51	0.05	23(04,49)	-0.44	Texas	0.56	0.02	11(04,29)	9.56
Louisiana	0.43	0.04	43(10,50)	-14.57	Utah	0.49	0.07	29(03,50)	-2.97
Maine	0.43	0.07	45(04,50)	-14.96	Vermont	0.45	0.11	40(01,50)	-10.49
Maryland	0.43	0.04	47(14,50)	-15.97 ^c	Virginia	0.50	0.03	26(06,47)	-1.63
Massachusetts	0.57	0.04	09(02,37)	13.02	Washington	0.50	0.04	24(05,47)	-0.54
Michigan	0.49	0.03	28(08,46)	-2.76	West Virginia	0.51	0.06	19(02,50)	0.26
Minnesota	0.51	0.04	21(04,48)	-0.13	Wisconsin	0.54	0.04	13(03,43)	6.78
Mississippi	0.43	0.05	46(07,50)	-15.90 ^c	Wyoming	-	-	-(,-)	-
Missouri	0.51	0.04	20(04,47)	-0.07					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.