Table 27.8 Cancer of the Urinary Bladder (Invasive and In Situ)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races				Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Female	
5-Year Relative Survival	(Percent)									
Year of Diagnosis:	(ICICCIIC)									
1960-1963 ^a	-		_	53	53	53	24	24	24	
1970-1973 ^a	-	-	-	61	61	60	36	38	24 27	
1975-1977 ^b	72.3	73.7	68.3	73.3	74.3	70.2	50.3	56.5	38.1	
1978-1980 ^b	73.9	75.1	70.4	74.7	75.4	72.3	54.5	62.6	40.1	
1981-1983 ^b	76.7	78.1	72.7	77.4	78.5	74.0	59.5	64.9	47.0	
1984-1986 ^b	76.7	78.1	72.6	77.2	78.5	73.6	59.4	62.3	52.8	
1987-1989 ^b	79.0	81.3	71.9	79.8	82.0	73.1	62.6	67.5	52.9	
1990-1992 ^b	79.2	81.7	71.9	80.2	82.6	73.1	63.1	65.7	58.5	
1993-1995 ^b	80.3	82.1	75.1	81.1	82.5	76.9	60.5	67.7	46.2	
1996-1998 ^b	79.1	80.4	75.4	79.8	81.1	75.9	62.7	65.6	56.0	
1999-2001 ^b	80.0	80.9	77.3	81.0	81.5	79.5	67.4	71.6	60.0	
2002-2005 ^b	79.9	81.7	74.6	80.8	82.4	75.9	63.2	67.4	55.6	
2006-2012 ^b	78.5 ^f	79.9 ^f	74.1 ^f	78.9 ^f	80.2 ^f	74.9 ^f	66.1 ^f	70.9 ^f	55.7	
5-Year Period Survival (Percent) ^{cd}									
2012	77.2	78.6	73.1	77.8	78.8	74.4	63.5	68.8	52.7	
Stage Distribution (%) 2 All Stages	006-2012 ^{ce}									
Number of cases	88,593	66,870	21,723	78,556	59,732	18,824	4,837	3,213	1,624	
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%	
In Situ	51	52	50	52	52	51	40	44	32	
Localized	34	35	33	34	35	33	37	36	39	
Regional	7	7	8	7	7	8	11	9	15	
Distant	4	4	6	4	4	5	8	7	9	
Unstaged	3	2	3	2	2	3	4	3	5	
5-Year Relative Survival Age at Diagnosis:	(Percent), 2	006-2012 ^c								
Ages <45	88.1	89.4	84.5	88.6	89.7	85.7	75.7	78.2	66.4	
Ages 45-54	84.4	85.2	82.2	85.5	85.8	84.4	69.6	73.7	59.0	
Ages 55-64	82.6	82.9	81.5	83.2	83.3	82.8	68.4	70.4	62.0	
Ages 65-74	79.5	80.4	76.3	80.2	80.9	77.3	65.0	69.2	57.1	
Ages 75+	70.4	72.7	64.3	70.9	72.9	65.3	54.6	60.9	46.2	
Ages <65	83.5	84.0	82.0	84.2	84.4	83.5	69.5	72.2	61.2	
Ages 65+	74.3	76.2	68.8	74.9	76.6	69.8	59.8	65.7	50.7	
Stage ^e :										
All Stages	77.5	78.9	73.0	78.0	79.2	74.1	63.9	68.8	53.9	
In Situ	95.9	95.6	96.7	96.1	95.7	97.0	89.9	90.4	87.8	
Localized	70.2	72.5	63.0	70.5	72.6	63.4	60.8	66.6	50.2	
	34.5	36.4	29.5	34.9	36.8	29.4	26.8		29.1	
Regional								24.2		
Distant	5.2	5.4	4.6	5.1	5.1	5.0	5.1	6.2	4.1	
Unstaged	45.9	50.5	35.5	44.5	48.9	34.0	38.0	48.8 ^g	21.5	
Based on End Results da SEER 9 areas (San Franc Based on follow-up of p	isco, Connect	icut, Detr								

SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Based on follow-up of patients into 2013.

Period survival provides a 2012 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival),

(2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival). Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding. The difference between 1975-1977 and 2006-2012 is statistically significant (p<.05).

The standard error is between 5 and 10 percentage points.

The standard error is greater than 10 percentage points.

Statistic could not be calculated due to fewer than 25 cases during the time period.

b

С

d

e

f

g

h

_