



Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	4.41	0.02		
High Five States				
Maine	5.99	0.26	01(01,04)	35.96 ^c
Vermont	5.60	0.39	02(01,25)	27.09 ^c
Nevada	5.28	0.21	03(01,19)	19.84 ^c
New Hampshire	5.13	0.26	04(01,29)	16.35 ^c
Ohio	5.11	0.09	05(02,14)	15.85 ^c
Low Five States				
Mississippi	3.71	0.16	47(33,50)	-15.79 ^c
Texas	3.71	0.06	48(42,50)	-15.83 ^c
South Dakota	3.70	0.27	49(21,50)	-16.04 ^c
Alaska	3.28	0.40	50(21,51)	-25.65 ^c
Hawaii	2.68	0.18	51(50,51)	-39.18 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.91	0.12	45(31,50)	-11.30	Montana	4.91	0.28	10(02,38)	11.34
Alaska	3.28	0.40	50(21,51)	-25.65 ^c	Nebraska	4.03	0.20	40(20,50)	-8.65
Arizona	4.07	0.11	37(25,48)	-7.67	Nevada	5.28	0.21	03(01,19)	19.84 ^c
Arkansas	4.56	0.17	21(06,38)	3.49	New Hampshire	5.13	0.26	04(01,29)	16.35 ^c
California	3.96	0.05	42(35,47)	-10.03	New Jersey	4.85	0.10	13(04,22)	10.00
Colorado	3.93	0.13	44(28,50)	-10.91	New Mexico	3.99	0.19	41(20,50)	-9.37
Connecticut	4.92	0.15	09(02,24)	11.65	New York	4.38	0.06	27(20,36)	-0.54
Delaware	4.62	0.30	20(02,47)	4.85	North Carolina	4.27	0.09	30(21,42)	-3.08
D.C.	3.85	0.37	46(08,51)	-12.68	North Dakota	4.22	0.32	34(03,50)	-4.26
Florida	4.42	0.06	26(19,34)	0.27	Ohio	5.11	0.09	05(02,14)	15.85 ^c
Georgia	4.45	0.10	24(14,36)	0.98	Oklahoma	4.53	0.15	23(08,38)	2.77
Hawaii	2.68	0.18	51(50,51)	-39.18 ^c	Oregon	4.76	0.15	17(04,30)	8.06
Idaho	4.21	0.23	35(08,49)	-4.38	Pennsylvania	4.84	0.08	15(05,22)	9.91
Illinois	4.34	0.08	28(20,38)	-1.42	Rhode Island	4.94	0.28	08(01,36)	12.18
Indiana	4.86	0.12	12(03,23)	10.30	South Carolina	4.25	0.13	32(19,45)	-3.56
Iowa	4.43	0.15	25(11,42)	0.51	South Dakota	3.70	0.27	49(21,50)	-16.04 ^c
Kansas	4.06	0.16	38(21,49)	-7.75	Tennessee	4.25	0.11	31(20,44)	-3.48
Kentucky	4.80	0.14	16(03,27)	8.98	Texas	3.71	0.06	48(42,50)	-15.83 ^c
Louisiana	4.16	0.14	36(20,47)	-5.61	Utah	3.95	0.20	43(21,50)	-10.35
Maine	5.99	0.26	01(01,04)	35.96 ^c	Vermont	5.60	0.39	02(01,25)	27.09 ^c
Maryland	4.99	0.13	07(02,21)	13.23	Virginia	4.23	0.10	33(21,44)	-4.02
Massachusetts	5.02	0.11	06(02,20)	13.90	Washington	4.88	0.12	11(03,22)	10.80
Michigan	4.85	0.09	14(04,22)	9.98	West Virginia	4.62	0.20	19(03,40)	4.90
Minnesota	4.04	0.12	39(25,48)	-8.25	Wisconsin	4.69	0.12	18(06,30)	6.32
Mississippi	3.71	0.16	47(33,50)	-15.79 ^c	Wyoming	4.55	0.40	22(02,50)	3.25
Missouri	4.28	0.11	29(19,43)	-2.79					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.