


Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2009-2013

Males

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	7.66	0.03		
High Five States				
Maine	10.52	0.54	01(01,05)	37.33 ^c
Vermont	9.41	0.79	02(01,39)	22.85 ^c
New Hampshire	9.24	0.55	03(01,25)	20.62 ^c
Ohio	8.93	0.18	04(02,15)	16.52 ^c
Maryland	8.92	0.28	05(02,19)	16.43 ^c
Low Five States				
South Dakota	6.50	0.56	47(16,51)	-15.16 ^c
Texas	6.43	0.12	48(41,49)	-16.02 ^c
D.C.	5.13	0.69	49(31,51)	-33.06 ^c
Hawaii	4.96	0.38	50(48,51)	-35.32 ^c
Alaska	4.77	0.72	51(34,51)	-37.71 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	7.12	0.27	36(20,48)	-7.10	Montana	8.77	0.58	07(01,37)	14.40
Alaska	4.77	0.72	51(34,51)	-37.71 ^c	Nebraska	6.98	0.40	42(17,49)	-8.96
Arizona	7.00	0.21	40(25,48)	-8.63	Nevada	8.78	0.41	06(01,27)	14.62
Arkansas	7.93	0.34	21(04,41)	3.45	New Hampshire	9.24	0.55	03(01,25)	20.62 ^c
California	6.81	0.10	44(34,47)	-11.16	New Jersey	8.49	0.21	13(03,23)	10.75
Colorado	6.72	0.27	45(27,49)	-12.26	New Mexico	6.98	0.39	41(17,49)	-8.85
Connecticut	8.53	0.32	11(02,27)	11.26	New York	7.75	0.13	22(17,33)	1.09
Delaware	7.21	0.59	35(04,49)	-5.90	North Carolina	7.45	0.20	29(19,42)	-2.83
D.C.	5.13	0.69	49(31,51)	-33.06 ^c	North Dakota	7.43	0.64	30(02,49)	-3.07
Florida	7.60	0.11	26(19,35)	-0.81	Ohio	8.93	0.18	04(02,15)	16.52 ^c
Georgia	7.93	0.23	20(08,34)	3.50	Oklahoma	7.73	0.31	23(09,43)	0.83
Hawaii	4.96	0.38	50(48,51)	-35.32 ^c	Oregon	8.32	0.30	18(03,30)	8.64
Idaho	7.27	0.46	34(08,48)	-5.09	Pennsylvania	8.64	0.16	08(03,19)	12.76
Illinois	7.53	0.17	27(18,39)	-1.69	Rhode Island	8.49	0.58	12(01,42)	10.86
Indiana	8.59	0.25	09(02,23)	12.10	South Carolina	7.29	0.28	31(17,47)	-4.85
Iowa	7.99	0.32	19(05,38)	4.33	South Dakota	6.50	0.56	47(16,51)	-15.16 ^c
Kansas	7.03	0.33	37(19,48)	-8.29	Tennessee	7.65	0.24	24(14,41)	-0.20
Kentucky	8.42	0.31	14(02,30)	9.88	Texas	6.43	0.12	48(41,49)	-16.02 ^c
Louisiana	7.02	0.28	39(21,48)	-8.37	Utah	6.84	0.40	43(19,49)	-10.79
Maine	10.52	0.54	01(01,05)	37.33 ^c	Vermont	9.41	0.79	02(01,39)	22.85 ^c
Maryland	8.92	0.28	05(02,19)	16.43 ^c	Virginia	7.28	0.22	32(20,45)	-4.98
Massachusetts	8.54	0.24	10(02,23)	11.48	Washington	8.37	0.24	16(03,26)	9.30
Michigan	8.36	0.19	17(04,24)	9.16	West Virginia	7.46	0.39	28(09,48)	-2.67
Minnesota	7.02	0.24	38(23,48)	-8.33	Wisconsin	8.40	0.25	15(03,27)	9.70
Mississippi	6.58	0.34	46(25,49)	-14.07	Wyoming	7.61	0.78	25(02,49)	-0.70
Missouri	7.28	0.23	33(20,46)	-5.02					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.